

Random attractors for rough differential equations

Luu Hoang Duc *

Abstract

We use the semigroup method to study the asymptotic behavior of rough differential equations, which consists of two steps of applying the continuous and discrete Gronwall lemmas. The existence of the global pullback attractor for the generated random dynamical system is then proved. We also derive an estimate for the diameter of the global attractor, and prove that for the linear diffusion function of linear form, the pullback attractor collapses to a random point.

Keywords: stochastic differential equations (SDE), rough path theory, rough differential equations, random dynamical systems, random attractors, exponential stability.

1 Introduction

This paper studies the asymptotic behavior of the rough differential equation

$$dy_t = [Ay_t + f(y_t)]dt + g(y_t)dx_t, t \in \mathbb{R}, y(0) = y_0 \in \mathbb{R}^d, \quad (1.1)$$

where we assume for simplicity that $A \in \mathbb{R}^{d \times d}$, $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$, $g : \mathbb{R}^d \rightarrow \mathbb{R}^{d \times m}$; f is globally Lipschitz continuous with Lipschitz coefficient C_f ; g either belongs to C_b^3 such that

$$C_g := \max \left\{ \|g\|_\infty, \|Dg\|_\infty, \|D_g^2\|_\infty, \|D_g^3\|_\infty \right\} < \infty, \quad (1.2)$$

or has a simple linear form $g(y) = Cy$, where $C \in \mathbb{R}^d \otimes \mathbb{R}^{d \times m}$. Such system is understood in the pathwise sense of a stochastic differential equation driven by a Hölder continuous stochastic process. Namely, we also assume that the driving path $x \in C^{\nu\text{-Hol}}(\mathbb{R}, \mathbb{R}^m) \subset C^{p\text{-var}}(\mathbb{R}, \mathbb{R}^m)$, with $\frac{1}{3} < \nu < \frac{1}{2}$, $p > \frac{1}{\nu}$ for simplicity, can be lifted into a realized component $\mathbf{x} = (x, \mathbb{X})$ of a stationary stochastic process $\mathbf{X}(\omega) = (1, x(\omega), \mathbb{X}(\omega))$, which has almost sure all realizations in the space $C^{\beta\text{-Hol}}(\mathbb{R}, T_1^2(\mathbb{R}^m)) \subset C^{p\text{-var}}(\mathbb{R}, T_1^2(\mathbb{R}^m))$, such that the estimate

$$E \left(\|x_{s,t}\|^p + \|\mathbb{X}_{s,t}\|^q \right) \leq C_{T,\nu} |t - s|^{p\nu}, \forall s, t \in [0, T]$$

holds for any $[0, T]$ for some constant $C_{T,\nu}$. In this circumstance, the solution is often solved in the sense of Friz-Victoir [12], and the existence and uniqueness theorem is recently proved in [23]. However, it is not clear how to apply the semigroup technique, which is well developed in [9] and [10] for Young differential equations [25], to estimate the rough path integrals. Therefore, we would like to study equation (1.1) in the sense of Gubinelli [16], in order to take advantage of the concept of rough integrals for controlled rough paths. Our aim is then to investigate the role of the driving noise in the longterm behavior of rough system (1.1).

Although no deterministic equilibrium such as the zero solution can in general be found, system (1.1) is expected to possess a pathwise attractor. The reader is referred to [13], [9], [10] and the

*Luu H. Duc is with Max-Planck-Institute for Mathematics in the Sciences, Leipzig, Germany, & Institute of Mathematics, Viet Nam Academy of Science and Technology duc.luu@mis.mpg.de, lhduc@math.ac.vn

references therein for recent development in studying the asymptotic behavior of Young differential equations, and [15], [7], [8] for rough differential equations.

The paper is organized as follows. Section 2 is devoted to present the existence, uniqueness and the norm estimates of the solution. In subsection 3.1, we introduce the generation of random dynamical system by the equation (1.1). Using Lemma 3.5, we prove the existence of a global random pullback attractor and estimate its diameter in Theorem 3.7 and Theorem 3.10. We also prove in Theorem 3.11 that in case $g(y) = Cy$, the attractor is actually a random point.

2 Rough differential equations

We would like to give a brief introduction to Young integrals. Given any compact time interval $I \subset \mathbb{R}$, let $C(I, \mathbb{R}^d)$ denote the space of all continuous paths $y : I \rightarrow \mathbb{R}^d$ equipped with sup norm $\|\cdot\|_{\infty, I}$ given by $\|y\|_{\infty, I} = \sup_{t \in I} \|y_t\|$, where $\|\cdot\|$ is the Euclidean norm in \mathbb{R}^d . We write $y_{s,t} := y_t - y_s$. For $p \geq 1$, denote by $\mathcal{C}^{p\text{-var}}(I, \mathbb{R}^d) \subset C(I, \mathbb{R}^d)$ the space of all continuous path $y : I \rightarrow \mathbb{R}^d$ which is of finite p -variation

$$\|y\|_{p\text{-var}, I} := \left(\sup_{\Pi(I)} \sum_{i=1}^n \|y_{t_i, t_{i+1}}\|^p \right)^{1/p} < \infty, \quad (2.1)$$

where the supremum is taken over the whole class of finite partition of I . $\mathcal{C}^{p\text{-var}}(I, \mathbb{R}^d)$ equipped with the p -var norm

$$\|y\|_{p\text{-var}, I} := \|y_{\min I}\| + \|y\|_{p\text{-var}, I},$$

is a nonseparable Banach space [12, Theorem 5.25, p. 92]. Also for each $0 < \alpha < 1$, we denote by $C^\alpha(I, \mathbb{R}^d)$ the space of Hölder continuous functions with exponent α on I equipped with the norm

$$\|y\|_{\alpha, I} := \|y_{\min I}\| + \|y\|_{\alpha, I} = \|y(a)\| + \sup_{s < t \in I} \frac{\|y_{s,t}\|}{(t-s)^\alpha},$$

A continuous map $\bar{\omega} : \Delta^2(I) \rightarrow \mathbb{R}^+$, $\Delta^2(I) := \{(s, t) : \min I \leq s \leq t \leq \max I\}$ is called a *control* if it is zero on the diagonal and superadditive, i.e. $\bar{\omega}_{t,t} = 0$ for all $t \in I$, and $\bar{\omega}_{s,u} + \bar{\omega}_{u,t} \leq \bar{\omega}_{s,t}$ for all $s \leq u \leq t$ in I .

Now, consider $y \in \mathcal{C}^{q\text{-var}}(I, \mathcal{L}(R^m, \mathbb{R}^d))$ and $x \in \mathcal{C}^{p\text{-var}}(I, \mathbb{R}^m)$ with $\frac{1}{p} + \frac{1}{q} > 1$, the Young integral $\int_I y_t dx_t$ can be defined as

$$\int_I y_s dx_s := \lim_{|\Pi| \rightarrow 0} \sum_{[u,v] \in \Pi} y_u x_{u,v},$$

where the limit is taken on all the finite partition $\Pi = \{\min I = t_0 < t_1 < \dots < t_n = \max I\}$ of I with $|\Pi| := \max_{[u,v] \in \Pi} |v - u|$ (see [25, p. 264-265]). This integral satisfies additive property by the construction, and the so-called Young-Loeve estimate [12, Theorem 6.8, p. 116]

$$\begin{aligned} \left\| \int_s^t y_u dx_u - y_s x_{s,t} \right\| &\leq K(p, q) \|y\|_{q\text{-var}, [s,t]} \|x\|_{p\text{-var}, [s,t]} \\ &\leq K(p, q) |t - s|^{\frac{1}{p} + \frac{1}{q}} \|y\|_{\frac{1}{p}, [s,t]} \|x\|_{\frac{1}{q}, [s,t]}, \end{aligned} \quad (2.2)$$

for all $[s, t] \subset I$, where

$$K(p, q) := (1 - 2^{1 - \frac{1}{p} - \frac{1}{q}})^{-1}. \quad (2.3)$$

We also introduce the construction of the integral using rough paths for the case $y, x \in C^\beta(I)$ when $\beta \in (\frac{1}{3}, \nu)$. To do that, we need to introduce the concept of rough paths. Following [11], a

couple $\mathbf{x} = (x, \mathbb{X})$, with $x \in C^\beta(I, \mathbb{R}^m)$ and $\mathbb{X} \in C_2^{2\beta}(\Delta^2(I), \mathbb{R}^m \otimes \mathbb{R}^m) := \{\mathbb{X} : \sup_{s < t} \frac{\|\mathbb{X}_{s,t}\|}{|t-s|^{2\beta}} < \infty\}$ where the tensor product $\mathbb{R}^m \otimes \mathbb{R}^n$ can be indentified with the matrix space $\mathbb{R}^{m \times n}$, is called a *rough path* if they satisfies Chen's relation

$$\mathbb{X}_{s,t} - \mathbb{X}_{s,u} - \mathbb{X}_{u,t} = x_{u,t} \otimes x_{s,u}, \quad \forall \min I \leq s \leq u \leq t \leq \max I. \quad (2.4)$$

\mathbb{X} is viewed as *postulating* the value of the quantity $\int_s^t x_{s,r} \otimes dx_r := \mathbb{X}_{s,t}$ where the right hand side is taken as a definition for the left hand side. Denote by $\mathcal{C}^\beta(I) \subset C^\beta \oplus C_2^{2\beta}$ the set of all rough paths in I , then \mathcal{C}^β is a closed set but not a linear space, equipped with the rough path semi-norm

$$\|\mathbf{x}\|_{\beta, I} := \|x\|_{\beta, I} + \|\mathbb{X}\|_{2\beta, \Delta^2(I)}^{\frac{1}{2}} < \infty. \quad (2.5)$$

Given fixed $\nu \in (\frac{1}{3}, \frac{1}{2})$, $\frac{1}{p} \in (\frac{1}{3}, \nu)$ and $\beta > \frac{1}{p}$, on each compact interval I such that $|I| = \max I - \min I \leq 1$, consider a rough path $\mathbf{x} = (x, \mathbb{X}) \in \mathcal{C}^{p\text{-var}}(I)$ with the p -var norm

$$\|\mathbf{x}\|_{p\text{-var}, I} := \left(\|x\|_{p\text{-var}, I}^p + \|\mathbb{X}\|_{q\text{-var}, I}^q \right)^{\frac{1}{p}}, \quad \text{where } q = \frac{p}{2}. \quad (2.6)$$

2.1 Controlled rough paths

Following [16], a path $y \in C^\beta(I, \mathcal{L}(\mathbb{R}^m, \mathbb{R}^d))$ is then called to be *controlled by* $x \in C^\beta(I, \mathbb{R}^m)$ if there exists a tube (y', R^y) with $y' \in C^\beta(I, \mathcal{L}(\mathbb{R}^m, \mathcal{L}(\mathbb{R}^m, \mathbb{R}^d)))$, $R^y \in C^{2\beta}(\Delta^2(I), \mathcal{L}(\mathbb{R}^m, \mathbb{R}^d))$ such that

$$y_{s,t} = y'_s \otimes x_{s,t} + R^y_{s,t}, \quad \forall \min I \leq s \leq t \leq \max I.$$

y' is called Gubinelli derivative of y , which is uniquely defined as long as $x \in C^\beta \setminus C^{2\beta}$ (see [11, Proposition 6.4]). The space $\mathcal{D}_x^{2\beta}(I)$ of all the couple (y, y') that is controlled by x will be a Banach space equipped with the norm

$$\begin{aligned} \|y, y'\|_{x, 2\beta, I} &:= \|y_{\min I}\| + \|y'_{\min I}\| + \| \|y, y'\| \|_{x, 2\beta, I}, \quad \text{where} \\ \| \|y, y'\| \|_{x, 2\beta, I} &:= \| \|y'\| \|_{\beta, I} + \| \|R^y\| \|_{2\beta, I}, \end{aligned}$$

where we omit the value space for simplicity of presentation. Now fix a rough path (x, \mathbb{X}) , then for any $(y, y') \in \mathcal{D}_x^{2\beta}(I)$, it can be proved that the function $F \in C^\beta(\Delta^2(I), \mathbb{R}^d)$ defined by

$$F_{s,t} := y_s \otimes x_{s,t} + y'_s \otimes \mathbb{X}_{s,t}$$

belongs to the space

$$\begin{aligned} C_2^{\beta, 3\beta}(I) &:= \left\{ F \in C^\beta(\Delta^2(I)) : F_{t,t} = 0 \quad \text{and} \right. \\ &\quad \left. \| \delta F \|_{3\beta, I} := \sup_{\min I \leq s \leq u \leq t \leq \max I} \frac{\|F_{s,t} - F_{s,u} - F_{u,t}\|}{|t-s|^{3\beta}} < \infty \right\}. \end{aligned}$$

Thanks to the sewing lemma [11, Lemma 4.2], the integral $\int_s^t y_u dx_u$ can be defined as

$$\int_s^t y_u dx_u := \lim_{|\Pi| \rightarrow 0} \sum_{[u,v] \in \Pi} [y_u \otimes x_{u,v} + y'_u \otimes \mathbb{X}_{u,v}]$$

where the limit is taken on all the finite partition Π of I with $|\Pi| := \max_{[u,v] \in \Pi} |v-u|$ (see [16]).

Moreover, there exists a constant $C_\beta = C_{\beta, |I|} > 1$ with $|I| := \max I - \min I$, such that

$$\left\| \int_s^t y_u dx_u - y_s x_{s,t} + y'_s \mathbb{X}_{s,t} \right\| \leq C_\beta |t-s|^{3\beta} \left(\|x\|_{\beta, [s,t]} \| \|R^y\| \|_{2\beta, \Delta^2[s,t]} + \| \|y'\| \|_{\beta, [s,t]} \| \mathbb{X} \|_{2\beta, \Delta^2[s,t]} \right). \quad (2.7)$$

From now on, if no other emphasis, we will simply write $\|x\|_\beta$ or $\|\mathbb{X}\|_{2\beta}$ without addressing the domain in I or $\Delta^2(I)$. In particular, for any $f \in C_b^3(\mathbb{R}^d, \mathbb{R}^d)$ we get the formula for integration by composition

$$f(x_t) = f(x_s) + \int_s^t \nabla f(x_u) dx_u + \frac{1}{2} \int_s^t \nabla^2 f(x_u) d[x]_{s,u},$$

where the last integral is understood in the Young sense and $[x]_{s,t} := x_{s,t} \otimes x_{s,t} - 2 \text{Sym}(\mathbb{X}_{s,t}) \in C^{2\beta}$. Notice that for geometric rough path $\mathbb{X}_{s,t} = \int_s^t x_{s,r} \otimes dx_r$, then $\text{Sym}(\mathbb{X}_{s,t}) = \frac{1}{2} x_{s,t} \otimes x_{s,t}$, thus $[x]_{s,t} \equiv 0$.

As proved in [16], the rough integral of controlled rough paths follows the rule of integration by parts. In practice, we would use the p -var norm

$$\begin{aligned} \|y, y'\|_{x,p,I} &:= \|y_{\min I}\| + \|y'_{\min I}\| + \|y, y'\|_{x,p,I}, \quad \text{where} \\ \|y, y'\|_{x,p,I} &:= \|y'\|_{p\text{-var},I} + \|R^y\|_{q\text{-var},I}. \end{aligned}$$

Thanks to the sewing lemma [5], we can use a similar version to (2.7) under p -var norm as follows.

$$\left\| \int_s^t y_u dx_u - y_s x_{s,t} + y'_s \mathbb{X}_{s,t} \right\| \leq C_p \left(\|x\|_{p\text{-var},[s,t]} \|R^y\|_{q\text{-var},\Delta^2[s,t]} + \|y'\|_{p\text{-var},[s,t]} \|\mathbb{X}\|_{q\text{-var},\Delta^2[s,t]} \right), \quad (2.8)$$

with constant $C_p > 1$ independent of \mathbf{x} and y .

2.2 Greedy times and integrability

In the following, we would like to construct a sequence of greedy times as presented in [4]. Given $\frac{1}{p} \in (\frac{1}{3}, \nu)$, we construct for any fixed $\gamma \in (0, 1)$ the sequence of greedy times $\{\tau_i(\gamma, I, p\text{-var})\}_{i \in \mathbb{N}}$ w.r.t. Hölder norms

$$\tau_0 = \min I, \quad \tau_{i+1} := \inf \left\{ t > \tau_i : \|\mathbf{x}\|_{p\text{-var},[\tau_i,t]} = \gamma \right\} \wedge \max I. \quad (2.9)$$

Denote by $N_{\gamma,I,p}(\mathbf{x}) := \sup\{i \in \mathbb{N} : \tau_i \leq \max I\}$. It follows that

$$N_{\gamma,I,p}(\mathbf{x}) \leq 1 + \gamma^{-p} \|\mathbf{x}\|_{p\text{-var},I}^p. \quad (2.10)$$

From now on, we would like to fix $\gamma = \frac{1}{4C_p C_g}$ and would like to write in short $N_{[a,b]}(\mathbf{x})$ for convenience.

2.3 Existence and uniqueness theorem

In this part, we would like to prove the existence and uniqueness theorem for rough differential equation (1.1), where the rough integral is understood in the sense of Gubinelli [16] for controlled rough paths. The idea is to prove first the existence, uniqueness and the differentiability w.r.t. the initial condition, of the solution of the rough differential equation

$$dy_t = g(y_t) dx_t, \quad \forall t \in [a, b], y_a \in \mathbb{R}^d, \quad (2.11)$$

and then to apply Doss-Sussmann technique [24] to transform the system to an equivalent ordinary differential equation. Note that the existence, uniqueness and continuity of the solution of (2.11) is already provided in [16], but the differentiability of the solution $y_t(\mathbf{x}, y_a)$ w.r.t. y_a is somehow missing due to the technical complex. We will derive below the proof for this statement.

Proposition 2.1 *The solution $y_t(\mathbf{x}, y_a)$ of (2.11) is uniformly continuous w.r.t. y_a , i.e.*

$$\begin{aligned}\|\bar{y} - y\|_{\infty, [a, b]} &\leq \|\bar{y}_a - y_a\| e^{(\log 2)\bar{N}_{[a, b]}(\mathbf{x})}, \\ \|\bar{y} - y, R\|_{p\text{-var}, [a, b]} &\leq \|\bar{y}_a - y_a\| \bar{N}_{[a, b]}^{\frac{p-1}{p}}(\mathbf{x}) e^{(\log 2)\bar{N}_{[a, b]}(\mathbf{x})} - \|\bar{y}_a - y_a\|,\end{aligned}\quad (2.12)$$

where $\bar{N}_{[a, b]}(\mathbf{x})$ is the maximal index of the maximal greedy time in the sequence

$$\tau_0 = a, \tau_{k+1} := \inf \left\{ t > \tau_k : \|\mathbf{x}\|_{p\text{-var}, [\tau_k, t]} = \left[8C_p C_g \left(1 + \frac{2}{C_p} N_{\frac{1}{8C_p C_g}, [a, b]}(\mathbf{x}) \right) \right]^{-1} \right\} \wedge b, \quad (2.13)$$

that lies in the interval $[a, b]$.

Proof: The proof is lengthy and is provided in the appendix. \square

Proposition 2.2 *The solution $y_t(\mathbf{x}, y_a)$ of (2.11) is differentiable w.r.t. initial condition y_a , moreover, its derivatives $\frac{\partial y_t}{\partial y_a}(\mathbf{x}, y_a)$ is the matrix solution of the linearized rough differential equation*

$$d\xi_t = Dg(y_t)\xi_t dx_t \quad (2.14)$$

Proof: The proof is lengthy and is provided in the appendix. \square

The following theorem shows a standard method to estimate the variation and the supremum norms of the solution of (1.1), by using Gronwall lemma and discretization scheme with the greedy times.

Theorem 2.3 *There exists a unique solution to (1.1) for any initial value, whose supremum and p -variation norms are estimated as follows*

$$\|y\|_{\infty, [a, b]} \leq \left[\|y_a\| + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_{[a, b]}(\mathbf{x}) \right] e^{4L(b-a)}, \quad (2.15)$$

$$\|y, R\|_{p\text{-var}, [a, b]} \leq \left[\|y_a\| + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_{[a, b]}(\mathbf{x}) \right] e^{4L(b-a)} N_{[a, b]}^{\frac{p-1}{p}}(\mathbf{x}) - \|y_a\|, \quad (2.16)$$

where $L = \|A\| + C_f$ and $\|y, R\|_{p\text{-var}, [s, t]} := \|y\|_{p\text{-var}, [s, t]} + \|R^y\|_{q\text{-var}, [s, t]}$.

Proof: Write in short $L = \|A\| + C_f$. The existence and uniqueness theorem follows [23] with the Doss-Sussmann method. Namely, using the integration by parts for the transformation $y_t = \varphi(t, x, z_t)$, it can be proved that there is a one-one corresponding between the solution of

$$dy_t = [Ay_t + f(y_t)]dt + g(y_t)dx_t = F(y_t)dt + g(y_t)dx_t. \quad (2.17)$$

and the solution of the ordinary differential equation

$$\dot{z}_t = \left[\frac{\partial \varphi}{\partial y}(t, x, z_t) \right]^{-1} F(\varphi(t, x, z_t)). \quad (2.18)$$

Since the right hand side of (2.18) satisfies the global Lipschitz continuity and linear growth, by similar arguments as in [23] there exists a unique solution given the initial value. That in turn proves the existence and uniqueness of system (2.17).

To prove (2.15), rewrite (1.1) in the integral form

$$y_{s, t} = \int_s^t [Ay_u + f(y_u)]du + \int_s^t g(y_u)dx_u. \quad (2.19)$$

Together with (1.2) and (2.8), we obtain

$$\begin{aligned}
& \|y_{s,t}\| \\
\leq & \int_s^t \left(\|Ay_u\| + \|f(y_u)\| \right) du + \left\| \int_s^t g(y_u) dx_u \right\| \\
\leq & \int_s^t (L\|y_u\| + \|f(0)\|) du + C_g \|x\|_{p\text{-var},[s,t]} + C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} + C_p \left\{ 2C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \|y\|_{p\text{-var},[s,t]} \right. \\
& \left. + \|x\|_{p\text{-var},[s,t]} \left[C_g \|R^y\|_{q\text{-var},[s,t]} + \frac{1}{2} C_g^2 \|x\|_{p\text{-var},[s,t]} \|y\|_{p\text{-var},[s,t]} \right] \right\} \\
\leq & \int_s^t (L\|y_u\| + \|f(0)\|) du + C_g \|x\|_{p\text{-var},[s,t]} + C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \\
& + C_p \left\{ \left[2C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} + \frac{1}{2} C_g^2 \|x\|_{p\text{-var},[s,t]}^2 \right] \vee C_g \|x\|_{p\text{-var},[s,t]} \right\} \left(\|y\|_{p\text{-var},[s,t]} + \|R^y\|_{q\text{-var},[s,t]} \right),
\end{aligned}$$

which yields

$$\begin{aligned}
& \|y\|_{p\text{-var},[s,t]} \\
\leq & \int_s^t (L\|y_u\| + \|f(0)\|) du + C_g \|x\|_{p\text{-var},[s,t]} + C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \\
& + C_p \left\{ \left[2C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} + \frac{1}{2} C_g^2 \|x\|_{p\text{-var},[s,t]}^2 \right] \vee C_g \|x\|_{p\text{-var},[s,t]} \right\} \left(\|y\|_{p\text{-var},[s,t]} + \|R^y\|_{q\text{-var},[s,t]} \right).
\end{aligned}$$

By similar arguments, we can show that

$$\begin{aligned}
& \|R^y\|_{q\text{-var},[s,t]} \\
\leq & \int_s^t (L\|y_u\| + \|f(0)\|) du + C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \\
& + C_p \left\{ \left[2C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} + \frac{1}{2} C_g^2 \|x\|_{p\text{-var},[s,t]}^2 \right] \vee C_g \|x\|_{p\text{-var},[s,t]} \right\} \left(\|y\|_{p\text{-var},[s,t]} + \|R^y\|_{q\text{-var},[s,t]} \right).
\end{aligned}$$

Therefore by assigning $\|y, R\|_{p\text{-var},[s,t]} = \|y\|_{p\text{-var},[s,t]} + \|R^y\|_{q\text{-var},[s,t]}$, we obtain

$$\begin{aligned}
\|y, R\|_{p\text{-var},[s,t]} & \leq 2 \int_s^t (L\|y\|_{p\text{-var},[s,u]} + L\|y_s\| + \|f(0)\|) du + C_g \|x\|_{p\text{-var},[s,t]} + 2C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \\
& + C_p \left\{ \left[2C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} + \frac{1}{2} C_g^2 \|x\|_{p\text{-var},[s,t]}^2 \right] \vee C_g \|x\|_{p\text{-var},[s,t]} \right\} \|y, R\|_{p\text{-var},[s,t]}.
\end{aligned} \tag{2.20}$$

Observe that if $2C_p C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} < 1$ then

$$2C_p C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} > C_p \left\{ \left[2C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} + \frac{1}{2} C_g^2 \|x\|_{p\text{-var},[s,t]}^2 \right] \vee C_g \|x\|_{p\text{-var},[s,t]} \right\}.$$

This follows that

$$\|y, R\|_{p\text{-var},[s,t]} \leq \int_s^t 4L \|y\|_{p\text{-var},[s,u]} du + 4(\|f(0)\| + L\|y_s\|)(t-s) + \frac{1}{C_p}$$

whenever $2C_p C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \leq \frac{1}{2}$. Applying the continuous Gronwall lemma 4.1, we obtain

$$\|y, R\|_{p\text{-var},[s,t]} \leq 4(\|f(0)\| + L\|y_s\|)(t-s) + \frac{1}{C_p}$$

$$\begin{aligned}
& + \int_s^t 4Le^{4L(t-u)} \left[4(\|f(0)\| + L\|y_s\|)(u-s) + \frac{1}{C_p} \right] du \\
& \leq \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} + \|y_s\| \right) e^{4L(t-s)} - \|y_s\|
\end{aligned} \tag{2.21}$$

whenever $4C_p C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \leq 1$. By constructing the sequence of greedy times $\{\tau_k(\frac{1}{4C_p C_g})\}_{k \in \mathbb{N}}$ on interval $[a, b]$, it follows from induction that

$$\begin{aligned}
\|y_{\tau_{k+1}}\| & \leq \|y\|_{\infty, [\tau_k, \tau_{k+1}]} \leq \|y\|_{p\text{-var}, [\tau_k, \tau_{k+1}]} \\
& \leq \left(\|y_{\tau_k}\| + \frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) e^{4L(\tau_{k+1} - \tau_k)} \\
& \leq e^{4L(\tau_{k+1} - \tau_0)} \|y_a\| + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) \sum_{j=0}^k e^{4L(\tau_{k+1} - \tau_j)} \\
& \leq \left[\|y_a\| + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) (k+1) \right] e^{4L(\tau_{k+1} - \tau_0)}, \quad \forall k = 0, \dots, N_{[a,b]}(\mathbf{x}) - 1,
\end{aligned}$$

which proves (2.15) since $\tau_{N_{[a,b]}(\mathbf{x})} = b$. On the other hand,

$$\begin{aligned}
\|y, R\|_{p\text{-var}, [\tau_k, \tau_{k+1}]} & \leq \|y_{\tau_k}\| \left(e^{4L(\tau_{k+1} - \tau_k)} - 1 \right) + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) e^{4L(\tau_{k+1} - \tau_k)} \\
& \leq \|y_a\| \left(e^{4L(\tau_{k+1} - \tau_0)} - e^{4L(\tau_k - \tau_0)} \right) + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) \left(e^{4L(\tau_{k+1} - \tau_0)} - e^{4L(\tau_k - \tau_0)} \right) \\
& \quad + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) e^{4L(\tau_{k+1} - \tau_k)}, \quad \forall k = 0, \dots, N_{[a,b]}(\mathbf{x}) - 1,
\end{aligned}$$

It then follows from inequality of p -variation seminorm in [8] that

$$\begin{aligned}
\|y, R\|_{p\text{-var}, [a,b]} & \leq N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x}) \sum_{k=0}^{N_{[a,b]}(\mathbf{x})-1} \|y, R\|_{q\text{-var}, [\tau_k, \tau_{k+1}]} \\
& \leq \|y_a\| N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x}) \sum_{k=0}^{N_{[a,b]}(\mathbf{x})-1} \left(e^{4L(\tau_{k+1} - \tau_0)} - e^{4L(\tau_k - \tau_0)} \right) \\
& \quad + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x}) \sum_{k=0}^{N_{[a,b]}(\mathbf{x})-1} \left(\sum_{j=0}^{k+1} e^{4L(\tau_{k+1} - \tau_j)} - \sum_{j=0}^k e^{4L(\tau_k - \tau_j)} \right) \\
& \leq N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x}) \left\{ \|y_a\| \left(e^{4L(b-a)} - 1 \right) + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) \left(\sum_{j=0}^{N_{[a,b]}(\mathbf{x})} e^{4L(b-\tau_j)} - 1 \right) \right\} \\
& \leq N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x}) \left[\|y_a\| + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_{[a,b]}(\mathbf{x}) \right] e^{4L(b-a)} - \|y_a\|
\end{aligned}$$

which proves (2.16). □

Following the same arguments line by line, we could prove similar estimates for $g = Cy$ as follows.

Theorem 2.4 *There exists a unique solution to the rough differential equation*

$$dy_t = [Ay_t + f(y_t)]dt + Cy_t dx_t, t \in \mathbb{R}, y(0) = y_0 \in \mathbb{R}^d, \tag{2.22}$$

for any initial value, whose supremum and p -variation norms are estimated as follows

$$\|y\|_{\infty,[a,b]} \leq \left[\|y_a\| + \frac{\|f(0)\|}{L} N_{[a,b]}(\mathbf{x}) \right] e^{4L(b-a) + \alpha N_{[a,b]}(\mathbf{x})}, \quad (2.23)$$

$$\|y, R\|_{p\text{-var},[a,b]} \leq \left[\|y_a\| + \frac{\|f(0)\|}{L} N_{[a,b]}(\mathbf{x}) \right] e^{4L(b-a) + \alpha N_{[a,b]}(\mathbf{x})} N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x}) - \|y_a\|, \quad (2.24)$$

where $L = \|A\| + C_f$, $\|y, R\|_{p\text{-var},[s,t]} := \|y\|_{p\text{-var},[s,t]} + \|R^y\|_{q\text{-var},[s,t]}$ and $\alpha = \log(1 + \frac{3}{2C_p})$.

Proof: Notice that the proof on the existence and uniqueness, as well as the Hölder norm estimates, of the solution of (2.4) is already given in [7]. To estimate the norms, we derive similar estimates as in the proof of Theorem 2.3 with note that

$$y'_s = C y_s, \quad [C y'_s] = C^2 y_s, \quad R_{s,t}^{C y} = C R_{s,t}^y.$$

Hence the estimate of $\|y, R\|_{p\text{-var},[s,t]}$ in (2.20) is of the form

$$\begin{aligned} & \|y, R\|_{p\text{-var},[s,t]} \\ & \leq 2 \int_s^t (L \|y\|_{p\text{-var},[s,u]} + L \|y_s\| + \|f(0)\|) du + \left(\|C\| \|x\|_{p\text{-var},[s,t]} + 2\|C\|^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \right) \|y_s\| \\ & \quad + 2C_p \left\{ \|C\|^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \vee \|C\| \|x\|_{p\text{-var},[s,t]} \right\} \|y, R\|_{p\text{-var},[s,t]}. \end{aligned}$$

As a result

$$\|y, R\|_{p\text{-var},[s,t]} \leq \int_s^t 4L \|y\|_{p\text{-var},[s,u]} du + 4(\|f(0)\| + L\|y_s\|)(t-s) + \frac{3}{2C_p} \|y_s\|$$

whenever $2C_p \|C\| \|\mathbf{x}\|_{p\text{-var},[s,t]} \leq \frac{1}{2}$. Applying the continuous Gronwall lemma 4.1, we obtain

$$\begin{aligned} \|y, R\|_{p\text{-var},[s,t]} & \leq 4(\|f(0)\| + L\|y_s\|)(t-s) + \frac{3}{2C_p} \|y_s\| \\ & \quad + \int_s^t 4L e^{4L(t-u)} \left[4(\|f(0)\| + L\|y_s\|)(u-s) + \frac{3}{2C_p} \|y_s\| \right] du \\ & \leq \left(\frac{\|f(0)\|}{L} + \left(1 + \frac{3}{2C_p}\right) \|y_s\| \right) e^{4L(t-s)} - \|y_s\| \end{aligned} \quad (2.25)$$

whenever $4C_p C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \leq 1$. The rest is the a direct consequence of [10, Theorem 2.1] (see also the proof of Propositions 2.1, 2.2 in the Appendix). \square

3 Random attractors

3.1 Generation of rough cocycle and rough flows

In this subsection we would like to present the generation of a random dynamical system from rough differential equations (1.1), which is based mainly on the work in [3] with only a small modification. Recall that a *random dynamical system* is defined by a mapping $\varphi(t, \omega)x_0 := x(t, \omega, x_0)$ on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ equipped with a metric dynamical system θ , i.e. $\theta_{t+s} = \theta_t \circ \theta_s$ for all $t, s \in \mathbb{R}$, such that $\varphi : \mathbb{R} \times \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a measurable mapping which is also continuous in t and x_0 and satisfies the cocycle property

$$\varphi(t+s, \omega) = \varphi(t, \theta_s \omega) \circ \varphi(s, \omega), \quad \forall t, s \in \mathbb{R},$$

(see [2]). Now denote by $T_1^2(\mathbb{R}^m) = 1 \oplus \mathbb{R}^m \oplus (\mathbb{R}^m \otimes \mathbb{R}^m)$ the set with the tensor product

$$(1, g^1, g^2) \otimes (1, h^1, h^2) = (1, g^1 + h^1, g^1 \otimes h^1 + g^2 + h^2), \quad \forall \mathbf{g} = (1, g^1, g^2), \mathbf{h} = (1, h^1, h^2) \in T_1^2(\mathbb{R}^m).$$

Then it can be shown that $(T_1^2(\mathbb{R}^m), \otimes)$ is a topological group with unit element $\mathbf{1} = (1, 0, 0)$.

For $\beta \in (\frac{1}{p}, \nu)$, denote by $\mathcal{C}^{0,p\text{-var}}([a, b], T_1^2(\mathbb{R}^m))$ the closure of $\mathcal{C}^\infty([a, b], T_1^2(\mathbb{R}^m))$ in $\mathcal{C}^{p\text{-var}}([a, b], T_1^2(\mathbb{R}^m))$, and by $\mathcal{C}_0^{0,p\text{-var}}(\mathbb{R}, T_1^2(\mathbb{R}^m))$ the space of all $x : \mathbb{R} \rightarrow \mathbb{R}^m$ such that $x|_I \in \mathcal{C}^{0,p\text{-var}}(I, T_1^2(\mathbb{R}^m))$ for each compact interval $I \subset \mathbb{R}$ containing 0. Then $\mathcal{C}_0^{0,p\text{-var}}(\mathbb{R}, T_1^2(\mathbb{R}^m))$ is equipped with the compact open topology given by the p -variation norm, i.e the topology generated by the metric:

$$d_p(\mathbf{x}_1, \mathbf{x}_2) := \sum_{k \geq 1} \frac{1}{2^k} (\|\mathbf{x}_1 - \mathbf{x}_2\|_{p\text{-var}, [-k, k]} \wedge 1),$$

where the p -var norm is given in (2.6). As a result, it is separable and thus a Polish space.

Let us consider a stochastic process $\bar{\mathbf{X}}$ defined on a probability space $(\bar{\Omega}, \bar{\mathcal{F}}, \bar{\mathbb{P}})$ with realizations in $(\mathcal{C}_0^{0,p\text{-var}}(\mathbb{R}, T_1^2(\mathbb{R}^m)), \mathcal{F})$. Assume further that $\bar{\mathbf{X}}$ has stationary increments. Assign

$$\Omega := \mathcal{C}_0^{0,p\text{-var}}(\mathbb{R}, T_1^2(\mathbb{R}^m))$$

and equip with the Borel σ -algebra \mathcal{F} and let \mathbb{P} be the law of $\bar{\mathbf{X}}$. Denote by θ the *Wiener-type shift*

$$(\theta_t \omega) = \omega_t^{-1} \otimes \omega_{t+\cdot}, \quad \forall t \in \mathbb{R}, \omega \in \mathcal{C}_0^{0,p\text{-var}}(\mathbb{R}, T_1^2(\mathbb{R}^m)), \quad (3.1)$$

and define the so-called *diagonal process* $\mathbf{X} : \mathbb{R} \times \Omega \rightarrow T_1^2(\mathbb{R}^m)$, $\mathbf{X}_t(\omega) = \omega_t$ for all $t \in \mathbb{R}, \omega \in \Omega$. Due to the stationarity of $\bar{\mathbf{X}}$, it can be proved that θ is invariant under \mathbb{P} , then forming a continuous (and thus measurable) dynamical system on $(\Omega, \mathcal{F}, \mathbb{P})$ [3, Theorem 5]. Moreover, \mathbf{X} forms a *p-rough path cocycle*, namely, $\mathbf{X}_\cdot(\omega) \in \mathcal{C}_0^{0,p\text{-var}}(\mathbb{R}, T_1^2(\mathbb{R}^m))$ for every $\omega \in \Omega$, which satisfies the *cocycle relation*:

$$\mathbf{X}_{t+s}(\omega) = \mathbf{X}_s(\omega) \otimes \mathbf{X}_t(\theta_s \omega), \quad \forall \omega \in \Omega, t, s \in \mathbb{R},$$

in the sense that $\mathbf{X}_{s,s+t} = \mathbf{X}_t(\theta_s \omega)$ with the increment notation $\mathbf{X}_{s,s+t} := \mathbf{X}_s^{-1} \otimes \mathbf{X}_{s+t}$. It is important to note that the two-parameter flow property

$$\mathbf{X}_{s,u} \otimes \mathbf{X}_{u,t} = \mathbf{X}_{s,t}, \quad \forall s, t \in \mathbb{R}$$

is equivalent to the fact that $\mathbf{X}_t(\omega) = (1, x_t(\omega), \mathbb{X}_{0,t}(\omega))$, where $x_\cdot(\omega) : \mathbb{R} \rightarrow \mathbb{R}^m$ and $\mathbb{X}_\cdot(\omega) : I \times I \rightarrow \mathbb{R}^m \otimes \mathbb{R}^m$ are random functions satisfying Chen's relation (2.4). To fulfill the Hölder continuity of almost all realizations, assume further that for any given $T > 0$, there exists a constant $C_{T,\nu}$ such that

$$E\left(\|x_{s,t}\|^p + \|\mathbb{X}_{s,t}\|^q\right) \leq C_{T,\nu} |t - s|^{\nu p}, \quad \forall s, t \in [0, T]. \quad (3.2)$$

Then due to the Kolmogorov criterion for rough paths [12, Appendix A.3], for any $\beta \in (\frac{1}{p}, \nu)$ there exists a version of ω -wise (x, \mathbb{X}) and random variables $K_\beta \in L^p, \mathbb{K}_\beta \in L^{\frac{p}{2}}$, such that, ω -wise speaking, for all $s, t \in I$,

$$\|x_{s,t}\| \leq K_\alpha |t - s|^\beta, \quad \|\mathbb{X}_{s,t}\| \leq \mathbb{K}_\beta |t - s|^{2\beta}, \quad \forall s, t \in \mathbb{R}$$

so that $(x, \mathbb{X}) \in \mathcal{C}^\beta$. Moreover, we could choose β such that

$$x \in \mathcal{C}^{0,\beta}(I) := \left\{ x \in \mathcal{C}^\beta : \lim_{\delta \rightarrow 0} \sup_{0 < t-s < \delta} \frac{\|x_{s,t}\|}{|t-s|^\beta} = 0 \right\},$$

$$\mathbb{X} \in \mathcal{C}^{0,2\beta}(\Delta^2(I)) := \left\{ \mathbb{X} \in \mathcal{C}^{2\beta}(\Delta^2(I)) : \lim_{\delta \rightarrow 0} \sup_{0 < t-s < \delta} \frac{\|\mathbb{X}_{s,t}\|}{|t-s|^{2\beta}} = 0 \right\},$$

then $C^{0,\beta}(I) \subset C^{0,\beta}(I) \oplus C^{0,2\beta}(\Delta^2(I))$ is separable due to the separability of $C^{0,\beta}(I)$ and $C^{0,2\beta}(\Delta^2(I))$. In particular, due to the fact that $\|\mathbf{X}_\cdot(\theta_h\omega)\|_{p\text{-var},[s,t]} = \|\mathbf{X}_\cdot(\omega)\|_{p\text{-var},[s+h,t+h]}$, it follows from Birkhoff ergodic theorem and (3.2) that

$$\Gamma(\mathbf{x}, p) := \limsup_{n \rightarrow \infty} \left(\frac{1}{n} \sum_{k=1}^n \|\theta_{-k}\mathbf{x}\|_{p\text{-var},[-1,1]}^p \right)^{\frac{1}{p}} = \left(E \|\mathbf{X}_\cdot(\cdot)\|_{p\text{-var},[-1,1]}^p \right)^{\frac{1}{p}} = \Gamma(p) \quad (3.3)$$

for almost all realizations \mathbf{x}_t of the form $\mathbf{X}_t(\omega)$. We assume additionally that $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ is ergodic.

Remark 3.1 It is important to note that, due to [3, Corollary 9], this construction is possible for $X : \mathbb{R} \rightarrow \mathbb{R}^m$ to be a continuous, centered Gaussian process with stationary increments and independent components, satisfying: there exists for any $T > 0$ a constant C_T such that for all $p \geq \frac{1}{\nu}$

$$E\|X_t - X_s\|^p \leq C_T |t - s|^{p\nu}, \quad \forall s, t \in [0, T]. \quad (3.4)$$

By Kolmogorov theorem, for any $\beta \in (\frac{1}{p}, \nu)$ and any interval $[0, T]$ almost all realization of X will be in $C^{0,\beta}([0, T])$. Then X has its covariance function with finite 2-dimensional ρ -variation on every square $[s, t]^2 \in \mathbb{R}^2$ for some $\rho \in [1, 2]$, and $\bar{\mathbf{X}}$ is the natural lift of X , in the sense of Friz-Victoir [12, Chapter 15], with sample paths in the space $\mathcal{C}_0^{0,\beta\text{-Hol}}(\mathbb{R}, T_1^2(\mathbb{R}^m))$, for every $p > 2\rho$.

For instance, such a stochastic process X , in particular, can be a m -dimensional fractional Brownian motion B^H with independent components [20] and Hurst exponent $H \in (\frac{1}{3}, \frac{1}{2})$, i.e. a family of $B^H = \{B_t^H\}_{t \in \mathbb{R}}$ with continuous sample paths and

$$E[B_t^H B_s^H] = \frac{1}{2} \left(t^{2H} + s^{2H} - |t - s|^{2H} \right) I^{m \times m}, \quad \forall t, s \in \mathbb{R}_+.$$

For any fixed interval $[0, T]$, the covariance of increments of fractional Brownian motions $R : [0, T]^4 \rightarrow \mathbb{R}^{m \times m}$, defined by

$$R \left(\begin{array}{cc} s & t \\ s' & t' \end{array} \right) := E(B_{s,t}^H B_{s',t'}^H)$$

is of finite ρ -variation norm for $\rho = \frac{1}{2H}$, i.e.

$$\|R\|_{I \times I', \rho} := \left\{ \sup_{\Pi(I), \Pi'(I')} \sum_{[s,t] \in I, [s',t'] \in I'} \left| R \left(\begin{array}{cc} s & t \\ s' & t' \end{array} \right) \right|^\rho \right\}^{\frac{1}{\rho}} < \infty,$$

and

$$\|R\|_{[s,t]^2, \rho} \leq M_{\rho, T} |t - s|^{\frac{1}{\rho}}, \quad \forall t, s \in [0, T].$$

Then one can prove that the integral in L^2 -sense

$$\mathbb{X}_{s,t}^{i,j} = \lim_{|\Pi| \rightarrow 0} \int_{\Pi} X_{s,r}^i dX_r^j = \lim_{|\Pi| \rightarrow 0} \sum_{[u,v] \in \Pi} X_{s,u}^i X_{u,v}^j, \quad \forall s, t \in [0, T]$$

is well-defined regardless of the chosen partition Π of $[s, t]$. Moreover,

$$\mathbb{X}_{s,t}^{i,i} = \frac{1}{2} (X_{s,t}^i)^2, \quad \mathbb{X}_{s,t}^{i,j} + \mathbb{X}_{s,t}^{j,i} = X_{s,t}^i X_{s,t}^j,$$

and for $\frac{1}{p} < \nu < \frac{1}{2\rho} = H$, there exist constants $C(p, \rho, m, T), C(p, \rho, m, T, \nu) > 0$ such that

$$\begin{aligned} E \left[\|X_{s,t}\|^p + \|\mathbb{X}_{s,t}\|^q \right] &\leq C(p, \rho, m, T) |t - s|^{\frac{p}{2\rho}} = C(p, \rho, m, T) |t - s|^{pH}, \quad \forall s, t \in [0, T], \\ E \left[\|\mathbb{X}\|_{\nu}^p + \|\mathbb{X}\|_{2\nu}^q \right] &\leq C(p, \rho, m, T, \nu) M^q. \end{aligned} \quad (3.5)$$

Therefore, almost sure all realizations $\mathbf{x} = (X, \mathbb{X})$ belong to the set $\mathcal{C}^\beta([0, T])$ and satisfy Chen's relation (2.4).

We reformulate a result from [3, Theorem 21] for our situation as follows.

Proposition 3.2 *Let $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ be a measurable metric dynamical system and let $\mathbf{X} : \mathbb{R} \times \Omega \rightarrow T_1^2(\mathbb{R}^m)$ be a p -rough cocycle for some $2 \leq p < 3$. Then there exists a unique continuous random dynamical system φ over $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ which solves the rough differential equation*

$$dy_t = [Ay_t + f(y_t)]dt + g(y_t)d\mathbf{X}_t(\omega), t \geq 0. \quad (3.6)$$

3.2 Existence of pullback attractors

Given a random dynamical system φ on \mathbb{R}^d , we follow [6], [2, Chapter 9] to present the notion of random pullback attractor. Recall that a set $\hat{M} = \{M(\omega)\}_{\omega \in \Omega}$ a *random set*, if $\omega \mapsto d(x|M(\omega))$ is \mathcal{F} -measurable for each $x \in \mathbb{R}^d$, where $d(E|F) = \sup\{\inf\{d(x,y)|y \in F\}|x \in E\}$ for E, F are nonempty subset of \mathbb{R}^d and $d(x|E) = d(\{x\}|E)$. An *universe* \mathcal{D} is a family of random sets which is closed w.r.t. inclusions (i.e. if $\hat{D}_1 \in \mathcal{D}$ and $\hat{D}_2 \subset \hat{D}_1$ then $\hat{D}_2 \in \mathcal{D}$). In our setting, we define the universe \mathcal{D} to be a family of random sets $D(\omega)$ which is *tempered* (see e.g. [2, pp. 164, 386]), namely $D(\omega)$ belongs to the ball $B(0, \rho(\omega))$ for all $\omega \in \Omega$ where the radius $\rho(\omega) > 0$ is a *tempered random variable*, i.e.

$$\lim_{t \rightarrow \pm\infty} \frac{1}{t} \log \rho(\theta_t \omega) = 0. \quad (3.7)$$

An invariant random compact set $\mathcal{A} \in \mathcal{D}$ is called a *pullback random attractor* in \mathcal{D} , if \mathcal{A} attracts any closed random set $\hat{D} \in \mathcal{D}$ in the pullback sense, i.e.

$$\lim_{t \rightarrow \infty} d(\varphi(t, \theta_{-t}\omega)\hat{D}(\theta_{-t}\omega)|\mathcal{A}(\omega)) = 0. \quad (3.8)$$

The existence of a random pullback attractor follows from the existence of a random pullback absorbing set (see [6, Theorem 3]). A random set $\mathcal{B} \in \mathcal{D}$ is called *pullback absorbing* in a universe \mathcal{D} if \mathcal{B} absorbs all sets in \mathcal{D} , i.e. for any $\hat{D} \in \mathcal{D}$, there exists a time $t_0 = t_0(\omega, \hat{D})$ such that

$$\varphi(t, \theta_{-t}\omega)\hat{D}(\theta_{-t}\omega) \subset \mathcal{B}(\omega), \text{ for all } t \geq t_0. \quad (3.9)$$

Given a universe \mathcal{D} and a random compact pullback absorbing set $\mathcal{B} \in \mathcal{D}$, there exists a unique random pullback attractor (which is then a weak attractor) in \mathcal{D} , given by

$$\mathcal{A}(\omega) = \bigcap_{s \geq 0} \overline{\bigcup_{t \geq s} \varphi(t, \theta_{-t}\omega)\mathcal{B}(\theta_{-t}\omega)}. \quad (3.10)$$

We need the following auxiliary results.

Proposition 3.3 *Assume that A has all eigenvalues of negative real parts. Then there exist constant $C_A \geq 1, \lambda_A > 0$ such that*

$$\|\Phi\|_{\infty, [a, b]} \leq C_A e^{-\lambda_A a}, \quad (3.11)$$

$$\|\Phi\|_{p\text{-var}, [a, b]} \leq \|A\| C_A e^{-\lambda_A a} (b - a), \quad \forall 0 \leq a < b, \quad (3.12)$$

where $\Phi(t) = e^{At}$.

Proof: See the proof in [10, Proposition 3.2]. □

Proposition 3.4 *Given (3.11) and (3.12), the following estimate holds: for any $0 \leq a < b \leq c$*

$$\begin{aligned} \left\| \int_a^b \Phi(c-s)g(y_s)dx_s \right\| &\leq C_A \left[1 + C_p \|A\| (b-a) \right] e^{-\lambda_A(c-b)} \left(C_g \|\mathbf{x}\|_{p\text{-var}, [a, b]} + C_g^2 \|\mathbf{x}\|_{p\text{-var}, [a, b]}^2 \right) \\ &\quad + 2C_p C_A e^{-\lambda_A(c-b)} \left\{ C_g^2 \|\mathbf{x}\|_{p\text{-var}, [a, b]}^2 \vee C_g \|\mathbf{x}\|_{p\text{-var}, [a, b]} \right\} \|y, R\|_{p\text{-var}, [a, b]}. \end{aligned} \quad (3.13)$$

Proof: Since

$$\Phi(c-t)g(y_t) - \Phi(c-s)g(y_s) = [\Phi(c-t) - \Phi(c-s)]g(y_t) + \Phi(c-s)\left([g(y)]'_s \otimes x_{s,t} + R_{s,t}^{g(y)}\right)$$

it follows that

$$\begin{aligned} [\Phi(c-\cdot)g(y)]'_s &= \Phi(c-s)[g(y)]'_s = \Phi(c-s)Dg(y_s)g(y_s), \\ \left\| R_{s,t}^{\Phi(c-\cdot)g(y)} \right\| &\leq \|\Phi(c-s)R_{s,t}^{g(y)}\| + \|\Phi(c-t) - \Phi(c-s)\| \|g(y_t)\|, \end{aligned}$$

which yields

$$\begin{aligned} \left\| [\Phi(c-\cdot)g(y)]' \right\|_{p\text{-var},[a,b]} &\leq 2C_g^2 \|\Phi(c-\cdot)\|_{\infty,[a,b]} \|y\|_{p\text{-var},[a,b]} + C_g^2 \|\Phi(c-\cdot)\|_{p\text{-var},[a,b]} \\ \left\| R^{\Phi(c-\cdot)g(y)} \right\|_{q\text{-var},[a,b]} &\leq \|\Phi(c-b)\| \left\| R^{g(y)} \right\|_{q\text{-var},[a,b]} + C_g \|\Phi(c-\cdot)\|_{q\text{-var},[a,b]}. \end{aligned}$$

Using (4.2) and (3.11), (3.12), we can now estimate

$$\begin{aligned} &\left\| \int_a^b \Phi(c-s)g(y_s)dx_s \right\| \\ &\leq \|\Phi(c-a)g(y_a)\| \|x_{a,b}\| + \|[\Phi(c-\cdot)g(y)]'_a\| \|\mathbb{X}_{a,b}\| \\ &\quad + C_p \left\{ \|x\|_{p\text{-var},[a,b]} \left\| R^{\Phi(c-\cdot)g(y)} \right\|_{q\text{-var},[a,b]} + \|\mathbb{X}\|_{q\text{-var},[a,b]} \left\| [\Phi(c-\cdot)g(y)]' \right\|_{p\text{-var},[a,b]} \right\} \\ &\leq C_A C_g e^{-\lambda_A(c-a)} \|x\|_{p\text{-var},[a,b]} + C_A C_g^2 e^{-\lambda_A(c-a)} \|\mathbb{X}\|_{q\text{-var},[a,b]} \\ &\quad + C_p \|\mathbb{X}\|_{q\text{-var},[a,b]} \left[2C_A C_g^2 e^{-\lambda_A(c-b)} \|y\|_{p\text{-var},[a,b]} + C_A C_g^2 \|A\| e^{-\lambda_A(c-b)}(b-a) \right] \\ &\quad + C_p \|x\|_{p\text{-var},[a,b]} \left\{ C_A C_g \|A\| e^{-\lambda_A(c-b)}(b-a) \right. \\ &\quad \left. + C_A e^{-\lambda_A(c-b)} \left(C_g \|R^y\|_{q\text{-var},[a,b]} + \frac{1}{2} C_g^2 \|x\|_{p\text{-var},[a,b]} \|y\|_{p\text{-var},[a,b]} \right) \right\} \\ &\leq C_A [1 + C_p \|A\| (b-a)] e^{-\lambda_A(c-b)} \left(C_g \|x\|_{p\text{-var},[a,b]} + C_g^2 \|\mathbb{X}\|_{q\text{-var},[a,b]} \right) \\ &\quad + C_p C_A e^{-\lambda_A(c-b)} \left\{ \left[2C_g^2 \|\mathbb{X}\|_{q\text{-var},[a,b]} + \frac{1}{2} C_g^2 \|x\|_{p\text{-var},[a,b]}^2 \right] \vee C_g \|x\|_{p\text{-var},[a,b]} \right\} \|y, R\|_{p\text{-var},[a,b]}. \end{aligned}$$

which, together proves (3.13). □

The following lemma is the crucial technique of this paper.

Lemma 3.5 *Assume that y_t satisfies*

$$y_t = \Phi(t)y_0 + \int_0^t \Phi(t-s)f(y_s)ds + \int_0^t \Phi(t-s)g(y_s)dx_s, \quad \forall t \geq 0. \quad (3.14)$$

Then for any $r > 0$ given and $n \geq 0$,

$$\begin{aligned} \|y_t\| e^{(\lambda_A - L_f)t} &\leq C_A \|y_0\| + \frac{C_A}{\lambda_A - L_f} \|f(0)\| \left(e^{(\lambda_A - L_f)t} - 1 \right) \\ &\quad + \sum_{k=0}^n e^{\lambda_A r} e^{(\lambda_A - L_f)kr} \left[\kappa_1(\mathbf{x}, \Delta_k^r) + \kappa_2(\mathbf{x}, \Delta_k^r) \|y, R\|_{p\text{-var}, \Delta_k^r} \right], \quad \forall t \in \Delta_n^r \end{aligned} \quad (3.15)$$

where $\Delta_k^r := [kr, (k+1)r]$, $L_f := C_A C_f$ and

$$\kappa_1(\mathbf{x}, [a, b]) := C_A [1 + C_p \|A\| (b-a)] \left(C_g \|\mathbf{x}\|_{p\text{-var},[a,b]} + C_g^2 \|\mathbf{x}\|_{p\text{-var},[a,b]}^2 \right) \quad (3.16)$$

$$\kappa_2(\mathbf{x}, [a, b]) := 2C_p C_A \left\{ C_g^2 \|\mathbf{x}\|_{p\text{-var},[a,b]}^2 \vee C_g \|\mathbf{x}\|_{p\text{-var},[a,b]} \right\}. \quad (3.17)$$

Proof: First, for any $t \in [nr, (n+1)r)$, it follows from (3.11) and the global Lipschitz continuity of f that

$$\begin{aligned} \|y_t\| &\leq \|\Phi(t)y_0\| + \int_0^t \|\Phi(t-s)f(y_s)\|ds + \left\| \int_0^t \Phi(t-s)g(y_s)dx_s \right\| \\ &\leq C_A e^{-\lambda_A t} \|y_0\| + \int_0^t C_A e^{-\lambda_A(t-s)} (C_f \|y_s\| + \|f(0)\|) ds + \left\| \int_0^t \Phi(t-s)g(y_s)dx_s \right\| \\ &\leq C_A e^{-\lambda_A t} \|y_0\| + \frac{C_A}{\lambda_A} \|f(0)\| (1 - e^{-\lambda_A t}) + \beta_t + C_A C_f \int_0^t e^{-\lambda_A(t-s)} \|y_s\| ds, \end{aligned}$$

where $\beta_t := \left\| \int_0^t \Phi(t-s)g(y_s)dx_s \right\|$. Multiplying both sides with $e^{\lambda_A t}$ yields

$$\|y_t\| e^{\lambda_A t} \leq C_A \|y_0\| + \frac{C_A}{\lambda_A} \|f(0)\| (e^{\lambda_A t} - 1) + \beta_t e^{\lambda_A t} + C_A C_f \int_0^t e^{\lambda_A s} \|y_s\| ds.$$

By applying the continuous Gronwall lemma 4.1, we obtain

$$\begin{aligned} \|y_t\| e^{\lambda_A t} &\leq C_A \|y_0\| + \frac{C_A}{\lambda_A} \|f(0)\| (e^{\lambda_A t} - 1) + \beta_t e^{\lambda_A t} \\ &\quad + \int_0^t L_f e^{L_f(t-s)} \left[C_A \|y_0\| + \frac{C_A}{\lambda_A} \|f(0)\| (e^{\lambda_A s} - 1) + \beta_s e^{\lambda_A s} \right] ds. \end{aligned}$$

Multiplying both sides with $e^{-L_f t}$ yields

$$\begin{aligned} \|y_t\| e^{(\lambda_A - L_f)t} &\leq C_A \|y_0\| e^{-L_f t} + \frac{C_A}{\lambda_A} \|f(0)\| \left(e^{(\lambda_A - L_f)t} - e^{-L_f t} \right) + \beta_t e^{(\lambda_A - L_f)t} \\ &\quad + \int_0^t L_f e^{-L_f s} \left[C_A \|y_0\| + \frac{C_A}{\lambda_A} \|f(0)\| (e^{\lambda_A s} - 1) + \beta_s e^{\lambda_A s} \right] ds \\ &\leq C_A \|y_0\| + \frac{C_A}{\lambda_A - L_f} \|f(0)\| \left(e^{(\lambda_A - L_f)t} - 1 \right) + \beta_t e^{(\lambda_A - L_f)t} + \int_0^t L_f \beta_s e^{(\lambda_A - L_f)s} ds. \end{aligned} \tag{3.18}$$

Next, observe from (3.13) that for all $s \leq t$

$$\begin{aligned} \beta_s e^{(\lambda_A - L_f)s} &= e^{(\lambda_A - L_f)s} \left\| \int_0^s \Phi(s-u)g(y_u)dx_u \right\| \\ &\leq e^{(\lambda_A - L_f)s} \sum_{k=0}^{\lfloor \frac{s}{r} \rfloor - 1} \left\| \int_{\Delta_k^r} \Phi(s-u)g(y_u)dx_u \right\| + \left\| \int_{r\lfloor s/r \rfloor}^s \Phi(s-u)g(y_u)dx_u \right\| \\ &\leq e^{(\lambda_A - L_f)s} \sum_{k=0}^{\lfloor \frac{s}{r} \rfloor - 1} e^{-\lambda_A(s-kr-r)} \left[\kappa_1(\mathbf{x}, \Delta_k^r) + \kappa_2(\mathbf{x}, \Delta_k^r) \|y, R\|_{p\text{-var}, \Delta_k^r} \right] \\ &\quad + e^{(\lambda_A - L_f)s} \left[\kappa_1(\mathbf{x}, [r\lfloor \frac{s}{r} \rfloor, s]) + \kappa_2(\mathbf{x}, [r\lfloor \frac{s}{r} \rfloor, s]) \|y, R\|_{p\text{-var}, [r\lfloor \frac{s}{r} \rfloor, s]} \right] \\ &\leq \sum_{k=0}^{\lfloor \frac{s}{r} \rfloor} e^{(\lambda_A - L_f)s} e^{-\lambda_A(s-kr-r)} \left[\kappa_1(\mathbf{x}, \Delta_k^r) + \kappa_2(\mathbf{x}, \Delta_k^r) \|y, R\|_{p\text{-var}, \Delta_k^r} \right] \\ &\leq e^{\lambda_A r} \sum_{k=0}^{\lfloor \frac{s}{r} \rfloor} e^{(\lambda_A - L_f)kr} e^{-L_f(s-kr)} \left[\kappa_1(\mathbf{x}, \Delta_k^r) + \kappa_2(\mathbf{x}, \Delta_k^r) \|y, R\|_{p\text{-var}, \Delta_k^r} \right]. \end{aligned} \tag{3.19}$$

Replacing (3.19) into (3.18) yields

$$\begin{aligned}
& \|y_t\| e^{(\lambda_A - L_f)t} \\
\leq & C_A \|y_0\| + \frac{C_A}{\lambda_A - L_f} \|f(0)\| \left(e^{(\lambda_A - L_f)t} - 1 \right) \\
& + e^{\lambda_A r} \sum_{k=0}^n e^{(\lambda_A - L_f)kr} e^{-L_f(t-kr)} \left[\kappa_1(\mathbf{x}, \Delta_k^r) + \kappa_2(\mathbf{x}, \Delta_k^r) \|y, R\|_{p\text{-var}, \Delta_k^r} \right] \\
& + e^{\lambda_A r} \int_0^t \sum_{k=0}^{\lfloor \frac{s}{r} \rfloor} e^{(\lambda_A - L_f)kr} e^{-L_f(s-kr)} \left[\kappa_1(\mathbf{x}, \Delta_k^r) + \kappa_2(\mathbf{x}, \Delta_k^r) \|y, R\|_{p\text{-var}, \Delta_k^r} \right] ds \\
\leq & C_A \|y_0\| + \frac{C_A}{\lambda_A - L_f} \|f(0)\| \left(e^{(\lambda_A - L_f)t} - 1 \right) \\
& + e^{\lambda_A r} \sum_{k=0}^n e^{(\lambda_A - L_f)kr} \left[\kappa_1(\mathbf{x}, \Delta_k^r) + \kappa_2(\mathbf{x}, \Delta_k^r) \|y, R\|_{p\text{-var}, \Delta_k^r} \right] \left(e^{-L_f(t-kr)} + \int_{kr}^t L_f e^{-L_f(s-kr)} ds \right) \\
\leq & C_A \|y_0\| + \frac{C_A}{\lambda_A - L_f} \|f(0)\| \left(e^{(\lambda_A - L_f)t} - 1 \right) \\
& + \sum_{k=0}^n e^{\lambda_A r} e^{(\lambda_A - L_f)kr} \left[\kappa_1(\mathbf{x}, \Delta_k^r) + \kappa_2(\mathbf{x}, \Delta_k^r) \|y, R\|_{p\text{-var}, \Delta_k^r} \right], \tag{3.20}
\end{aligned}$$

where we use the fact that $e^{-L_f(t-kr)} + \int_{kr}^t L_f e^{-L_f(s-kr)} ds = 1$ for all $t \geq kr$. The continuity of y at $t = (n+1)r$ then proves (3.15). \square

We need one more auxiliary proposition.

Proposition 3.6 *Define*

$$G(\mathbf{x}, [a, b]) := e^{\lambda_A + 4L(b-a)} \kappa_2(\mathbf{x}, [a, b]) N_{[a, b]}^{\frac{p-1}{p}}(\mathbf{x}), \tag{3.21}$$

$$\begin{aligned}
H(\mathbf{x}, [a, b]) & := e^{\lambda_A r} \left\{ \frac{C_A \|f(0)\| (e^\lambda - 1)}{\lambda} + \kappa_1(\mathbf{x}, [a, b]) \right. \\
& \quad \left. + \kappa_2(\mathbf{x}, [a, b]) \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) e^{4L(b-a)} N_{[a, b]}^{\frac{2p-1}{p}}(\mathbf{x}) \right\}, \tag{3.22}
\end{aligned}$$

and

$$b(\mathbf{x}) := \sum_{k=1}^{\infty} e^{-\lambda k} H(\theta_{-k}\mathbf{x}, [-1, 1]) \prod_{j=1}^{k-1} \left[1 + G(\theta_{-j}\mathbf{x}, [-1, 1]) \right] \tag{3.23}$$

(which can be infinity), where $\lambda := \lambda_A - C_A C_f$, κ_1, κ_2 are given by (3.16), (3.17). Assume further that

$$\lambda > \hat{G} := \frac{1}{2} C_A e^{\lambda_A + 8L} \left\{ \left[4C_p C_g \Gamma(p) \right]^p + \left[4C_p C_g \Gamma(p) \right] \right\}. \tag{3.24}$$

Then $b(\mathbf{x})$ is finite and tempered a.s., i.e.

$$\lim_{t \rightarrow \pm\infty} \frac{1}{t} \log b(\theta_t \mathbf{x}) = 0. \tag{3.25}$$

Proof: Assign $\Delta_k = [k, k+1]$ and $N_k(\mathbf{x}) := N_{\Delta_k}(\mathbf{x})$. Observe from (2.10) that

$$\begin{aligned}
N_k^{\frac{p-1}{p}}(\mathbf{x}) & \leq \left(1 + [4C_p C_g]^p \| \mathbf{x} \|_{p\text{-var}, \Delta_k}^p \right)^{\frac{p-1}{p}} \leq 1 + [4C_p C_g]^{p-1} \| \mathbf{x} \|_{p\text{-var}, \Delta_k}^{p-1}, \\
N_k^{\frac{2p-1}{p}}(\mathbf{x}) & \leq \left(1 + [4C_p C_g]^p \| \mathbf{x} \|_{p\text{-var}, \Delta_k}^p \right)^{\frac{2p-1}{p}} \leq 2^{\frac{p-1}{p}} \left(1 + [4C_p C_g]^{2p-1} \| \mathbf{x} \|_{p\text{-var}, \Delta_k}^{2p-1} \right),
\end{aligned}$$

In addition, it follows from (3.24) that $2C_g\Gamma(p) < 2C_pC_g\Gamma(p) < 1$. As a result, a direct computation shows that

$$G(\mathbf{x}, [a, b]) \leq \frac{1}{2}C_A e^{\lambda_A + 4L(b-a)} \left\{ 4C_pC_g \|\mathbf{x}\|_{p\text{-var}, [a, b]} + [4C_pC_g]^p \|\mathbf{x}\|_{p\text{-var}, [a, b]}^p \right\} \left\{ C_g \|\mathbf{x}\|_{p\text{-var}, [a, b]} \vee 1 \right\}. \quad (3.26)$$

Due to the inequality $\log(1 + \frac{v}{4C_p}(u + u^p)(u \vee 1)) \leq v(u + u^p)$ for all $u \geq 0, v \geq \frac{1}{2}$, (3.26) yields

$$\log \left(1 + G(\mathbf{x}, [-1, 1]) \right) \leq \frac{1}{2}C_A e^{\lambda_A + 8L} \left\{ 4C_pC_g \|\mathbf{x}\|_{p\text{-var}, [-1, 1]} + [4C_pC_g]^p \|\mathbf{x}\|_{p\text{-var}, [-1, 1]}^p \right\}.$$

It follows that for a.s. all x ,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \prod_{k=0}^{n-1} \left[1 + G(\theta_{-k}\mathbf{x}, [-1, 1]) \right] &= \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \log \left[1 + G(\theta_k\mathbf{x}, [-1, 1]) \right] \\ &\leq \frac{1}{2}C_A e^{\lambda_A + 8L} \left\{ \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \|\theta_{-k}\mathbf{x}\|_{p\text{-var}, [-1, 1]}^p + \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \|\theta_{-k}\mathbf{x}\|_{p\text{-var}, [-1, 1]} \right\} \\ &\leq \frac{1}{2}C_A e^{\lambda_A + 8L} \left\{ [4C_pC_g\Gamma(p)]^p + [4C_pC_g\Gamma(p)] \right\} = \hat{G}. \end{aligned}$$

Similarly, it is easy to show from (3.22) and (3.16), (3.17) that $E|\log H(\mathbf{x}, [-1, 1])| < \infty$, thus

$$\limsup_{n \rightarrow \infty} \frac{\log H(\theta_n\mathbf{x}, [-1, 1])}{n} = \limsup_{n \rightarrow \infty} \frac{\log H(\theta_{-n}\mathbf{x}, [-1, 1])}{n} = 0.$$

Hence, there exists for each $0 < 2\delta < \lambda - \hat{G}$ an $n_0 = n_0(\delta, x)$ such that for all $n \geq n_0$,

$$e^{(-\delta + \hat{G})n} \leq \prod_{k=0}^{n-1} \left[1 + G(\theta_{-k}\mathbf{x}, [-1, 1]) \right], \quad \prod_{k=0}^{n-1} \left[1 + G(\theta_k\mathbf{x}, [-1, 1]) \right] \leq e^{(\delta + \hat{G})n}$$

and

$$e^{-\delta n} \leq H(\theta_{-n}\mathbf{x}, [-1, 1]), \quad H(\theta_n\mathbf{x}, [-1, 1]) \leq e^{\delta n}.$$

Consequently,

$$\begin{aligned} b(\mathbf{x}) &\leq \sum_{k=1}^{n_0-1} e^{-\lambda k} H(\theta_{-k}\mathbf{x}, [-1, 1]) \prod_{j=1}^{k-1} \left(1 + M_1 C_g G(\theta_{-j}\mathbf{x}, [-1, 1]) \right) + \sum_{k=n_0}^{\infty} e^{-(\lambda - 2\delta - \hat{G})k} \\ &\leq \sum_{k=1}^{n_0-1} e^{-\lambda k} H(\theta_{-k}\mathbf{x}, [-1, 1]) \prod_{j=1}^{k-1} \left(1 + M_1 C_g G(\theta_{-j}\mathbf{x}, [-1, 1]) \right) + \frac{e^{-(\lambda - 2\delta - \hat{G})n_0}}{1 - e^{-(\lambda - 2\delta - \hat{G})}} \end{aligned}$$

which is finite. The proof on the temperedness of $b(\mathbf{x})$ is proved similarly to [10, Appendix]. \square

We are now able to formulate the first main result of the paper.

Theorem 3.7 *Assume that A has all eigenvalues of negative real parts with λ_A satisfying (3.11) and (3.12), and f is globally Lipschitz continuous such that $\lambda_A > C_f C_A$. Assume further that the driving path x satisfies (3.3). Then under the condition*

$$\lambda_A - C_A C_f > \frac{1}{2}C_A e^{\lambda_A + 8(\|A\| + C_f)} \left\{ [4C_pC_g\Gamma(p)]^p + [4C_pC_g\Gamma(p)] \right\}, \quad (3.27)$$

where $\Gamma(p) = \left(E \|\mathbf{Z}\|_{p\text{-var}, [-1, 1]}^p \right)^{\frac{1}{p}}$, the random dynamical system φ possesses a pullback attractor $\mathcal{A}(\mathbf{x})$.

Proof: Using the rule of integration by parts for rough integrals, it is easy to prove that y_t satisfies

$$y_t = \Phi(t)y_0 + \int_0^t \Phi(t-s)f(y_s)ds + \int_0^t \Phi(t-s)g(y_s)dx_s. \quad (3.28)$$

Then by applying Proposition 3.5 and using the estimate in (2.15)

$$\|y, R\|_{p\text{-var}, \Delta_k} \leq \|y_k\| e^{4L} N_k^{\frac{p-1}{p}}(\mathbf{x}) + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) e^{4L} N_k^{\frac{2p-1}{p}}(\mathbf{x}),$$

where $N_k(x) := N_{\Delta_k}(x)$, we obtain

$$\begin{aligned} \|y_n\| e^{\lambda n} &\leq C_A \|y_0\| + (e^{\lambda n} - 1) \frac{C_A \|f(0)\|}{\lambda} \\ &\quad + e^{\lambda A} \sum_{k=0}^{n-1} e^{\lambda k} \left\{ \kappa_1(\mathbf{x}, \Delta_k) + \kappa_2(\mathbf{x}, \Delta_k) \left[\|y_k\| e^{4L} N_k^{\frac{p-1}{p}}(\mathbf{x}) + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) e^{4L} N_k^{\frac{2p-1}{p}}(\mathbf{x}) \right] \right\} \\ &\leq C_A \|y_0\| + \frac{C_A \|f(0)\| (e^\lambda - 1)}{\lambda} \sum_{k=0}^{n-1} e^{\lambda k} + \sum_{k=0}^{n-1} e^{\lambda k} \kappa_2(\mathbf{x}, \Delta_k) e^{4L} N_k^{\frac{p-1}{p}}(\mathbf{x}) \|y_k\| \\ &\quad + e^{\lambda A} \sum_{k=0}^{n-1} e^{\lambda k} \left\{ \kappa_1(\mathbf{x}, \Delta_k) + \kappa_2(\mathbf{x}, \Delta_k) \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) e^{4L} N_k^{\frac{2p-1}{p}}(\mathbf{x}) \right\} \\ &\leq C_A \|y_0\| + \sum_{k=0}^{n-1} \kappa_2(\mathbf{x}, \Delta_k) e^{4L} N_k^{\frac{p-1}{p}}(\mathbf{x}) e^{\lambda k} \|y_k\| \\ &\quad + e^{\lambda A} \sum_{k=0}^{n-1} e^{\lambda k} \left\{ \frac{C_A \|f(0)\| (e^\lambda - 1)}{\lambda} + \kappa_1(\mathbf{x}, \Delta_k) + \kappa_2(\mathbf{x}, \Delta_k) \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) e^{4L} N_k^{\frac{2p-1}{p}}(\mathbf{x}) \right\}. \end{aligned} \quad (3.29)$$

By assigning $a := C_A \|y_0\|$, $u_k := \|y_k\| e^{\lambda k}$, $k \geq 0$ and using (3.21), (3.22), we obtain

$$u_n \leq a + \sum_{k=0}^{n-1} G(x, \Delta_k) u_k + \sum_{k=0}^{n-1} e^{\lambda k} H(x, \Delta_k). \quad (3.30)$$

We are now in the position to apply Lemma 4.2, so that

$$\|y_n(\mathbf{x}, y_0)\| \leq C_A \|y_0\| e^{-\lambda n} \prod_{k=0}^{n-1} [1 + G(\theta_k x, [0, 1])] + \sum_{k=0}^{n-1} e^{-\lambda(n-k)} H(\theta_k x, [0, 1]) \prod_{j=k+1}^{n-1} [1 + G(\theta_j x, [0, 1])]. \quad (3.31)$$

Now using (2.15), it follows that for any $t \in [n, n+1]$

$$\begin{aligned} \|y_t(\mathbf{x}, y_0)\| &\leq \left[\|y_n(\mathbf{x}, y_0)\| + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_n(\mathbf{x}) \right] e^{4L} \\ &\leq C_A e^{4L} \|y_0\| e^{-\lambda n} \prod_{k=0}^{n-1} [1 + G(\theta_k \mathbf{x}, [0, 1])] \\ &\quad + \sum_{k=0}^{n-1} e^{-\lambda(n-k)} e^{4L} H(\theta_k \mathbf{x}, [0, 1]) \prod_{j=k+1}^{n-1} [1 + G(\theta_j \mathbf{x}, [0, 1])] + e^{4L} \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_n(\mathbf{x}). \end{aligned} \quad (3.32)$$

Consequently, by assigning x with $\theta_{-t}x$ in (3.32), we obtain

$$\|y_t(\theta_{-t}\mathbf{x}, y_0(\theta_{-t}x))$$

$$\begin{aligned}
&\leq C_A e^{4L} \|y_0\| e^{-\lambda n} \prod_{k=0}^{n-1} \left[1 + G(\theta_{k-t}\mathbf{x}, [0, 1]) \right] \\
&\quad + \sum_{k=0}^{n-1} e^{-\lambda(n-k)} e^{4L} H(\theta_{k-t}\mathbf{x}, [0, 1]) \prod_{j=k+1}^{n-1} \left[1 + G(\theta_{j-t}\mathbf{x}, [0, 1]) \right] + e^{4L} \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_n(\theta_{-t}\mathbf{x}) \\
&\leq C_A e^{4L} \|y_0\| e^{-\lambda n} \prod_{k=0}^{n-1} \left[1 + G(\theta_{k-n}\mathbf{x}, [-1, 1]) \right] + e^{4L} \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_{[-1,1]}(\mathbf{x}) \\
&\quad + \sum_{k=0}^{n-1} e^{-\lambda(n-k)} e^{4L} H(\theta_{k-n}\mathbf{x}, [-1, 1]) \prod_{j=k+1}^{n-1} \left[1 + G(\theta_{j-n}\mathbf{x}, [-1, 1]) \right] \tag{3.33}
\end{aligned}$$

We are now in the position to apply Proposition 3.6 into (3.33) so that for $t \in \Delta_n$ with $0 < \delta < \frac{1}{2}(\lambda - \hat{G})$ and n large enough

$$\begin{aligned}
\|y_t(\theta_{-t}\mathbf{x}, y_0)\| &\leq C_A e^{4L} \|y_0(\theta_{-t}\mathbf{x})\| \exp \left\{ -(\lambda - \hat{G} - \delta) n \right\} + e^{4L} \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_{[-1,1]}(\mathbf{x}) \\
&\quad + \sum_{k=1}^n e^{-\lambda k} e^{4L} H(\theta_{-k}\mathbf{x}, [-1, 1]) \prod_{j=k+1}^{n-1} \left[1 + G(\theta_{-j}\mathbf{x}, [-1, 1]) \right] \\
&\leq C_A e^{4L} \|y_0(\theta_{-t}\mathbf{x})\| \exp \left\{ -(\lambda - \hat{G} - \delta) n \right\} + b(\mathbf{x}) + \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_{[-1,1]}(\mathbf{x}), \tag{3.34}
\end{aligned}$$

where $b(\mathbf{x})$ is given by (3.23). This implies that, starting from any point $y_0(\theta_{-t}\mathbf{x}) \in D(\theta_{-t}\mathbf{x})$ which is tempered due to (3.7), there exists n large enough such that for $t \in [n, n+1]$

$$\|y_t(\theta_{-t}\mathbf{x}, y_0)\| \leq 1 + b(\mathbf{x}) + e^{4L} \left(\frac{\|f(0)\|}{L} + \frac{1}{C_p} \right) N_{[-1,1]}(\mathbf{x}) =: \hat{b}(\mathbf{x}). \tag{3.35}$$

Moreover, the temperedness of $\hat{b}(\mathbf{x})$ follows directly from the temperedness (3.25) of $b(\mathbf{x})$ and of $\|\mathbf{x}\|_{p\text{-var}, [-1,1]}$. Therefore, there exists a compact absorbing set $\mathcal{B}(\mathbf{x}) = \bar{B}(0, \hat{b}(\mathbf{x}))$ and thus a pullback attractor $\mathcal{A}(\mathbf{x})$ for system (1.1) which is given by (3.10). \square

Remark 3.8 (i), Assume that $f(0) = g(0) = 0$ so that $y \equiv 0$ is a solution of (1.1). Then (3.27) in Theorem 3.7 is the exponential stability criterion for the trivial attractor $\mathcal{A}(\mathbf{x}) \equiv 0$.

(ii) It is important to note that the term $e^{\lambda_A + 8(\|A\| + C_f)}$ in (3.27) is the unavoidable effect from the discretization scheme.

(iii), A similar proof of Theorem 3.7 using step size r with $\Delta_k = [kr, (k+1)r]$ then leads to a criterion for the existence of a global random pullback attractor

$$\lambda_A - C_A C_f > \frac{1}{2r} C_A e^{\left[\lambda_A + 8(\|A\| + C_f) \right] r} \left\{ \left[4C_p C_g \Gamma(p, r) \right]^p + \left[4C_p C_g \Gamma(p, r) \right] \right\} \tag{3.36}$$

where $\Gamma(p, r) = \left(E \|Z\|_{p\text{-var}, [-r, r]}^p \right)^{\frac{1}{p}}$ for almost sure all realizations x . As a result, the final criterion can be optimized to

$$\lambda_A - C_A C_f > \inf_{r>0} \frac{1}{2r} C_A e^{\left[\lambda_A + 8(\|A\| + C_f) \right] r} \left\{ \left[4C_p C_g \Gamma(p, r) \right]^p + \left[4C_p C_g \Gamma(p, r) \right] \right\}.$$

In the following, we are going to prove that the diameter of the random attractor can be controlled by parameter C_g . We first introduce a quantity.

Proposition 3.9 *Assume that x satisfies (3.3). Then under criterion (3.27), the following quantity is well defined and finite*

$$\begin{aligned} \xi(\mathbf{x}) &:= e^{4L} \sum_{k=1}^{\infty} \left(\|\theta_{-k}\mathbf{x}\|_{p\text{-var},[0,1]} + \|\theta_{-k}\mathbf{x}\|_{p\text{-var},[0,1]}^2 \right) \times \\ &\quad \times e^{-\lambda(n-k)} \left[\max \left\{ \frac{\|f(0)\|}{L}, \frac{1}{C_p} \right\} N_{[0,1]}(\theta_{-k}\mathbf{x}) + \hat{b}(\theta_{-k}\mathbf{x}) \right] N_{[0,1]}(\theta_{-k}\mathbf{x})^{\frac{p-1}{p}}. \end{aligned} \quad (3.37)$$

Proof: Observe that the existence of $\Gamma(\mathbf{x}, p)$ implies the temperedness of $N_{[0,1]}(\mathbf{x}), N_{[0,1]}(\mathbf{x})^{\frac{p-1}{p}}$ and $\|\mathbf{x}\|_{p\text{-var},[0,1]} + \|\mathbf{x}\|_{p\text{-var},[0,1]}^2$. The convergence of the series in (3.37) can then be proved similarly to the convergence of $\hat{b}(x)$ in Proposition 3.6. \square

Theorem 3.10 *Under the assumptions of Theorem 3.7, the diameter of \mathcal{A} is estimated as*

$$\text{diam}(\mathcal{A}(\mathbf{x})) \leq 2C_p C_A (1 + \|A\|) (C_g \vee C_g^2) \xi(\mathbf{x}) \quad (3.38)$$

where $\xi(\mathbf{x})$ is given in (3.37).

Proof: The existence of the pullback attractor \mathcal{A} is followed by Theorem 3.7. Take any two points $a_1, a_2 \in \mathcal{A}(x)$. For a given $n \in \mathbb{N}$, assign $x^* := \theta_{-n}x$ and consider the equation

$$dy_t = [Ay_t + f(y_t)]dt + g(y_t)dx_t^*. \quad (3.39)$$

Due to the invariance of \mathcal{A} under the flow, there exist $b_1, b_2 \in \mathcal{A}(x^*)$ such that $a_i = y_n(x^*, b_i)$. Put $z_t = z_t(x^*) := y_t(x^*, b_1) - y_t(x^*, b_2)$ then $z_n(x^*) = a_1 - a_2$ and we have

$$dz_t = [Az_t + P(t, z_t)]dt + Q(t, z_t)dx_t^* \quad (3.40)$$

where we write in short $y_t^1 = y_t(x^*, b_1)$ and

$$\begin{aligned} P(t, z_t) &= f(y(t, x^*, b_2)) - f(y(t, x^*, b_1)) = f(y_t^1 + z_t) - f(y_t^1), \\ Q(t, z_t) &= g(y(t, x^*, b_2)) - g(y(t, x^*, b_1)) = g(y_t^1 + z_t) - g(y_t^1). \end{aligned}$$

Observe that

$$\|P(t, z) - P(t, z')\| \leq C_f \|z - z'\|, \quad \|Q(t, z) - Q(t, z')\| \leq C_g \|z - z'\|$$

and $P(t, 0) = Q(t, 0) \equiv 0$. Consequently,

$$\|P(t, z_t)\| \leq C_f \|z_t\|, \quad \|Q(t, z_t)\| \leq C_g \|z_t\|.$$

Using the rule of integration by parts for rough integrals, it is easy to see that z_t satisfies the equation

$$z_t = z_a + \int_a^t \Phi(t-s)P(s, z_s)ds + \int_a^t \Phi(t-s)Q(s, z_s)dx_s^*.$$

On the other hand, a direct computation shows that

$$\begin{aligned} [\Phi(c-\cdot)Q(\cdot, z)]'_s &= \Phi(c-s)[g(y^1)]'_s - \Phi(c-s)[g(y^s)]'_s \\ &= \Phi(c-s)Dg(y_s^1)g(y_s^1) - \Phi(c-s)Dg(y_s^2)g(y_s^2) \\ \left\| R_{s,t}^{\Phi(c-\cdot)Q(\cdot, z)} \right\| &\leq \|\Phi(c-s)R_{s,t}^Q\| + \|\Phi(c-t) - \Phi(c-s)\| \|Q(t, z_t)\| \end{aligned} \quad (3.41)$$

which yields

$$\begin{aligned}
\|[\Phi(c-\cdot)Q(\cdot, z)]'\|_{p\text{-var}, [a, b]} &\leq 2C_g^2 \|\Phi(c-\cdot)\|_{\infty, [a, b]} \left(\|y^1\|_{p\text{-var}, [a, b]} + \|y^2\|_{p\text{-var}, [a, b]} \right) \\
&\quad + 2C_g^2 \|\Phi(c-\cdot)\|_{p\text{-var}, [a, b]} \\
\|R^{\Phi(c-\cdot)Q(\cdot, z)}\|_{q\text{-var}, [a, b]} &\leq \|\Phi(c-b)\| \left(\|R^{g(y^1)}\|_{q\text{-var}, [a, b]} + \|R^{g(y^2)}\|_{q\text{-var}, [a, b]} \right) \\
&\quad + 2C_g \|\Phi(c-\cdot)\|_{q\text{-var}, [a, b]}
\end{aligned} \tag{3.42}$$

As a result, we use similar estimate to Proposition 3.4 to obtain

$$\begin{aligned}
&\left\| \int_a^b \Phi(c-s)Q(s, z_s) dx_s \right\| \\
&\leq \|\Phi(c-a)Q(a, z_a)\| \|x\|_{p\text{-var}, [a, b]} + \|[\Phi(c-\cdot)Q(\cdot, z)]'_a\| \|\mathbb{X}\|_{q\text{-var}, [a, b]} \\
&\quad + C_p \left\{ \|x\|_{p\text{-var}, [a, b]} \|R^{\Phi(c-\cdot)Q}\|_{q\text{-var}, [a, b]} + \|\mathbb{X}\|_{q\text{-var}, [a, b]} \|[\Phi(c-\cdot)Q]'\|_{p\text{-var}, [a, b]} \right\} \\
&\leq C_A e^{-\lambda_A(c-a)} \left(C_g \|x\|_{p\text{-var}, [a, b]} + 2C_g^2 \|\mathbb{X}\|_{q\text{-var}, [a, b]} \right) \|z_a\| \\
&\quad + C_A C_p \|A\| (b-a) e^{-\lambda_A(c-b)} \left(C_g \|x\|_{p\text{-var}, [a, b]} + C_g^2 \|\mathbb{X}\|_{q\text{-var}, [a, b]} \right) \\
&\quad + C_p C_A e^{-\lambda_A(c-b)} \left\{ \left[2C_g^2 \|\mathbb{X}\|_{q\text{-var}, [a, b]} + \frac{1}{2} C_g^2 \|x\|_{p\text{-var}, [a, b]}^2 \right] \vee C_g \|x\|_{p\text{-var}, [a, b]} \right\} \times \\
&\quad \times \left(\|y^1, R^{y^1}\|_{p\text{-var}, [a, b]} + \|y^2, R^{y^2}\|_{p\text{-var}, [a, b]} \right).
\end{aligned} \tag{3.43}$$

Now, repeating the estimate in the proof of Theorem 3.7 with $\beta_t^* = \|\int_0^t \Phi(t-s)Q(s, z_s) dx_s^*\|$ we obtain

$$e^{\lambda_A t} \|z_t\| \leq C_A \|z_0\| + e^{\lambda_A t} \beta_t^* + L_f \int_0^t \left(C_A \|z_0\| + e^{\lambda_A s} \beta_s^* \right) e^{L_f(t-s)} ds$$

and then

$$e^{\lambda t} \|z_t\| \leq C_A \|z_0\| + e^{\lambda t} \beta_t^* + L_f \int_0^t e^{\lambda s} \beta_s^* ds \tag{3.44}$$

Similarly to (3.19) we have

$$\begin{aligned}
\beta_t^* e^{\lambda t} &= e^{\lambda t} \left\| \int_0^t \Phi(t-s)Q(s, z_s) dx_s^* \right\| \\
&\leq e^{\lambda t} \sum_{k=0}^{\lfloor t \rfloor} C_p C_A (1 + \|A\|) \left(C_g \|\mathbf{x}^*\|_{p\text{-var}, \Delta_k} + C_g^2 \|\mathbf{x}^*\|_{p\text{-var}, \Delta_k}^2 \right) e^{-\lambda_A(t-k-1)} \times \\
&\quad \times \left(\|y_k^1\| + \|y_k^2\| + \|y^1, R^{y^1}\|_{p\text{-var}, \Delta_k} + \|y^2, R^{y^2}\|_{p\text{-var}, \Delta_k} \right).
\end{aligned} \tag{3.45}$$

Therefore the similar estimate to (3.20) in Lemma 3.5 shows that

$$\begin{aligned}
e^{\lambda t} \|z_t\| &\leq C_A \|z_0\| + C_p C_A (1 + \|A\|) \sum_{k=0}^{\lfloor t \rfloor} \left(C_g \|\mathbf{x}^*\|_{p\text{-var}, \Delta_k} + C_g^2 \|\mathbf{x}^*\|_{p\text{-var}, \Delta_k}^2 \right) \times \\
&\quad \times e^{\lambda k} \left(\|y_k^1\| + \|y_k^2\| + \|y^1, R^{y^1}\|_{p\text{-var}, \Delta_k} + \|y^2, R^{y^2}\|_{p\text{-var}, \Delta_k} \right).
\end{aligned} \tag{3.46}$$

Since $b_i \in \mathcal{A}(\mathbf{x}^*)$ for $i = 1, 2$, it follows from the invariance of \mathcal{A} that $y^i(k, \mathbf{x}^*, b_i) \in \mathcal{A}(\theta_k \mathbf{x}^*)$. Moreover, it follows from (3.10) and (3.35) that

$$\sup_{y \in \mathcal{A}(\mathbf{x})} \|y\| \leq \hat{b}(\mathbf{x}). \quad (3.47)$$

Indeed, taking $y^* \in \mathcal{A}(\mathbf{x})$ be arbitrary, it follows from (3.10) that there exists a sequence $t_k \rightarrow \infty$ such that

$$y^* = \lim_k \varphi(t_k, \theta_{-t_k} \mathbf{x}, y_0(\theta_{-t_k} \mathbf{x}))$$

where $y_0(\theta_{-t_k} \mathbf{x}) \in \mathcal{B}(\theta_{-t_k} \mathbf{x})$. Since $\hat{b}(\mathbf{x})$ is tempered, by choosing t_k large enough so that (3.35) holds, we conclude that (3.47) holds. As a consequence, (3.47) yields $\|y^1(k, \mathbf{x}^*, b_1)\| \leq \hat{b}(\theta_k \mathbf{x}^*)$. Similarly, $\|z_0\| \leq \|b_1\| + \|b_2\| < 2\hat{b}(\mathbf{x}^*)$. On the other hand, due to (2.16) and (3.46) yields

$$\begin{aligned} \|z_n\| &\leq 2C_A \hat{b}(\mathbf{x}^*) e^{-\lambda n} + 2C_p C_A (1 + \|A\|) (C_g \vee C_g^2) \sum_{k=0}^{n-1} \left(\|\mathbf{x}^*\|_{p\text{-var}, \Delta_k} + \|\mathbf{x}^*\|_{p\text{-var}, \Delta_k}^2 \right) e^{-\lambda(n-k)} \times \\ &\quad \times \left[\max \left\{ \frac{\|f(0)\|}{L}, \frac{1}{C_p} \right\} N_{\Delta_k}(\mathbf{x}^*) + \hat{b}(\theta_k \mathbf{x}^*) \right] e^{4L} N_{\Delta_k}(\mathbf{x}^*)^{\frac{p-1}{p}} \\ &\leq 2C_A \hat{b}(\theta_{-n} \mathbf{x}) e^{-\lambda n} + 2C_p C_A (1 + \|A\|) (C_g \vee C_g^2) \sum_{k=1}^n \left(\|\theta_{-k} \mathbf{x}\|_{p\text{-var}, [0,1]} + \|\theta_{-k} \mathbf{x}\|_{p\text{-var}, [0,1]}^2 \right) \times \\ &\quad \times e^{-\lambda(n-k)} \left[\max \left\{ \frac{\|f(0)\|}{L}, \frac{1}{C_p} \right\} N_{[0,1]}(\theta_{-k} \mathbf{x}) + \hat{b}(\theta_{-k} \mathbf{x}) \right] e^{4L} N_{[0,1]}(\theta_{-k} \mathbf{x})^{\frac{p-1}{p}}. \end{aligned} \quad (3.48)$$

Letting n tend to infinity, the first term in the last line of (3.48) tends to zero due to the temperedness of $\hat{b}(\mathbf{x})$. Hence it follows from (3.37) in Proposition 3.9 that

$$\|a_1 - a_2\| \leq 2C_p C_A (1 + \|A\|) (C_g \vee C_g^2) \xi(\mathbf{x})$$

which proves (3.38). □

In the rest of the paper, we are going to prove the result on one-point attractor in case g is of linear form, as proved in [10] for Young equations.

Theorem 3.11 *Assume that $g(y) = Cy$ is a linear map. Then under the condition*

$$\lambda_A - C_A C_f > 2C_A [1 + \|A\|] e^{\lambda_A + 4(\|A\| + C_f)} \left\{ \left[4C_p \|C\| \Gamma(p) \right] + \left[4C_p \|C\| \Gamma(p) \right]^p \right\}, \quad (3.49)$$

the attractor is a random point, i.e. $\mathcal{A}(\mathbf{x}) = \{a(\mathbf{x})\}$.

Proof: With the setting in the proof of Theorem 3.10 for $g(y) = Cy$, observe that $Q(t, z_t) = Cz_t$ and

$$\begin{aligned} \|\Phi(c - \cdot) Cz_t\|_{p\text{-var}, [a,b]} &\leq \|\Phi(c - \cdot)\|_{\infty, [a,b]} \|C\|^2 \|z\|_{p\text{-var}, [a,b]} + \|\Phi(c - \cdot)\|_{p\text{-var}, [a,b]} \|C\|^2 \|z\|_{\infty, [a,b]} \\ \|\mathbb{R}^{\Phi(c - \cdot)} Cz_t\|_{q\text{-var}, [a,b]} &\leq \|\Phi(c - \cdot)\|_{\infty, [a,b]} \|C\| \|\mathbb{R}^z\|_{q\text{-var}, [a,b]} + \|\Phi(c - \cdot)\|_{q\text{-var}, [a,b]} \|C\| \|z\|_{\infty, [a,b]}. \end{aligned}$$

As a result, the estimate in (3.43) is of the form

$$\begin{aligned} &\left\| \int_a^b \Phi(c - s) Q(s, z_s) dx_s^* \right\| \\ &\leq \|\Phi(c - a) Cz_a\| \|x^*\|_{p\text{-var}, [a,b]} + \|\Phi(c - a) C^2 z_a\| \|\mathbb{X}^*\|_{q\text{-var}, [a,b]} \end{aligned}$$

$$\begin{aligned}
& +C_p \left\{ \|x^*\|_{p\text{-var},[a,b]} \left\| R^{\Phi(c-\cdot)} C \right\|_{q\text{-var},[a,b]} + \|\mathbb{X}^*\|_{q\text{-var},[a,b]} \left\| [\Phi(c-\cdot)C]' \right\|_{p\text{-var},[a,b]} \right\} \\
& \leq C_p C_A [1 + \|A\|(b-a)] e^{-\lambda A(c-b)} \left(\|C\| \|\mathbf{x}^*\|_{p\text{-var},[a,b]} + \|C\|^2 \|\mathbf{x}^*\|_{p\text{-var},[a,b]}^2 \right) \times \\
& \quad \times \left(\|z_a\| + \|z, R^z\|_{p\text{-var},[a,b]} \right). \tag{3.50}
\end{aligned}$$

Meanwhile, similar estimates to (2.24) in Theorem 2.4, with $P(t, 0) = 0$, show that

$$\begin{aligned}
\|z, R^z\|_{p\text{-var},[a,b]} + \|z_a\| & \leq \|z_a\| e^{4L(b-a) + \alpha N_{[a,b]}(\mathbf{x}^*)} N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x}^*), \quad \text{with} \\
N_{[a,b]}(\mathbf{x}^*) & \leq 1 + [4C_p \|C\|]^p \|\mathbf{x}^*\|_{p\text{-var},[a,b]}^p.
\end{aligned}$$

As a result, (3.46) has the form

$$\begin{aligned}
e^{\lambda n} \|z_n\| & \leq C_A \|z_0\| + C_p C_A [1 + \|A\|] e^{\lambda A} \sum_{k=0}^{n-1} \left(\|C\| \|\mathbf{x}^*\|_{p\text{-var},\Delta_k} + \|C\|^2 \|\mathbf{x}^*\|_{p\text{-var},\Delta_k}^2 \right) \times \\
& \quad \times e^{4L + \alpha N_{\Delta_k}(\mathbf{x}^*)} N_{\Delta_k}^{\frac{p-1}{p}}(\mathbf{x}^*) e^{\lambda k} \|z_k\| \\
& \leq C_A \|z_0\| + \sum_{k=0}^{n-1} I_{\Delta_k}(\mathbf{x}^*) e^{\lambda k} \|z_k\|,
\end{aligned}$$

where

$$I_{[a,b]}(\mathbf{x}) = C_p C_A [1 + \|A\|] e^{\lambda A} \left(\|C\| \|\mathbf{x}\|_{p\text{-var},[a,b]} + \|C\|^2 \|\mathbf{x}\|_{p\text{-var},[a,b]}^2 \right) e^{4L + \alpha N_{[a,b]}(\mathbf{x})} N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x})$$

is the function of \mathbf{x} . Now applying the discrete Gronwall lemma, we obtain

$$e^{\lambda n} \|z_n\| \leq C_A \|z_0\| \prod_{k=0}^{n-1} \left[1 + I_{[0,1]}(\theta_{k-n} \mathbf{x}) \right]$$

Hence, it follows from Birkhorff's ergodic theorem that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \|z_n\| \leq -\lambda + \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \log \left[1 + I_{[0,a]}(\theta_{-k} \mathbf{x}) \right] \leq -\lambda + E \log \left[1 + I_{[0,1]}(\mathbf{x}) \right].$$

Given C_p and α , it follows from the estimate of $N_{[0,1]}(\mathbf{x})$ and the inequalities

$$\begin{aligned}
\log(1 + ue^v) & \leq v + \log(1 + u), \quad \forall u, v \geq 0, \\
\log \left[1 + \frac{(2C_p + 3)v}{4C_p} (1 + u^{p-1})(u + u^2) \right] & \leq (2 - \alpha)v(u + u^p), \quad \forall u \geq 0, v \geq 1,
\end{aligned}$$

that

$$\begin{aligned}
& \log \left[1 + I_{[0,1]}(\mathbf{x}) \right] \\
& \leq \log \left\{ 1 + C_p C_A [1 + \|A\|] e^{\lambda A} e^{4L + \alpha} N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x}) \left(\|C\| \|\mathbf{x}\|_{p\text{-var},[a,b]} + \|C\|^2 \|\mathbf{x}\|_{p\text{-var},[a,b]}^2 \right) \right\} \\
& \quad + \alpha \left(4C_p \|C\| \right)^p \|\mathbf{x}\|_{p\text{-var},[0,1]}^p \\
& \leq \log \left\{ 1 + \frac{2C_p + 3}{4C_p} C_A [1 + \|A\|] e^{\lambda A + 4L} \left[1 + (4C_p \|C\|)^{p-1} \|\mathbf{x}\|_{p\text{-var},[0,1]}^{p-1} \right] \right\} \times \\
& \quad \times \left(4C_p \|C\| \|\mathbf{x}\|_{p\text{-var},[a,b]} + \left[4C_p \|C\| \right]^2 \|\mathbf{x}\|_{p\text{-var},[a,b]}^2 \right) \left\} + \alpha \left(4C_p \|C\| \right)^p \|\mathbf{x}\|_{p\text{-var},[0,1]}^p
\end{aligned}$$

$$\leq 2C_A[1 + \|A\|]e^{\lambda_A+4L} \left\{ \left[4C_p \|C\| \|\mathbf{x}\|_{p\text{-var},[0,1]} \right] + \left[4C_p \|C\| \|\mathbf{x}\|_{p\text{-var},[0,1]} \right]^p \right\}$$

Therefore we finally obtain

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \|z_n\| \leq -\lambda + 2C_A[1 + \|A\|]e^{\lambda_A+4L} \left\{ \left[4C_p \|C\| \Gamma(p) \right] + \left[4C_p \|C\| \Gamma(p) \right]^p \right\} < 0$$

under the condition (3.49). This follows that $\lim_{n \rightarrow \infty} \|a_1 - a_2\| = 0$ or \mathcal{A} is an one point set. \square

4 Appendix

Lemma 4.1 (Continuous Gronwall Lemma) *Assume that $u_t, \alpha_t, \beta > 0$ such that*

$$u_t \leq \alpha_t + \int_a^t \beta u_s ds, \forall t \geq a.$$

Then

$$u_t \leq \alpha_t + \int_a^t \beta e^{\beta(t-s)} \alpha_s ds, \forall t \geq a.$$

Proof: See [1, Lemma 6.1, p 89]. \square

Lemma 4.2 (Discrete Gronwall Lemma) *Let a be a non negative constant and u_n, α_n, β_n be nonnegative sequences satisfying*

$$u_n \leq a + \sum_{k=0}^{n-1} \alpha_k u_k + \sum_{k=0}^{n-1} \beta_k, \quad \forall n \geq 1$$

then

$$u_n \leq \max\{a, u_0\} \prod_{k=0}^{n-1} (1 + \alpha_k) + \sum_{k=0}^{n-1} \beta_k \prod_{j=k+1}^{n-1} (1 + \alpha_j) \quad (4.1)$$

for all $n \geq 1$.

Proof: See [10, Appendix 4.2] \square

Proof: [**Proposition 2.1**] The proof is divided into several steps.

Step 1: First we would like to estimate the solution norms of (2.4). To do that, observe that

$$\left\| \int_s^t g(y_u) dx_u - g(y_s) x_{s,t} - [g(y)]'_s \mathbb{X}_{s,t} \right\| \leq C_\beta (t-s)^{3\beta} \left[\|x\|_{\beta, [s,t]} \left\| R^{g(y)} \right\|_{2\beta, [s,t]} + \|\mathbb{X}\|_{2\beta, [s,t]} \left\| [g(y)]' \right\|_{\beta, [s,t]} \right]$$

due to (2.7). It then follows that y is controlled by x with $y' = g(y)$. Since

$$\begin{aligned} g(y_t) - g(y_s) &= \int_0^1 Dg(y_s + \eta y_{s,t}) y_{s,t} d\eta \\ &= Dg(y_s) y'_s \otimes x_{s,t} + \int_0^1 Dg(y_s + \eta y_{s,t}) R_{s,t}^y d\eta + \int_0^1 [Dg(y_s + \eta y_{s,t}) - Dg(y_s)] y'_{s,t} \otimes x_{s,t} d\eta, \end{aligned}$$

it easy to show that $[g(y)]'_s = Dg(y_s)g(y_s)$, where we use (1.2) to estimate

$$\begin{aligned} \|R_{s,t}^{g(y)}\| &\leq \int_0^1 \|Dg(y_s + \eta y_{s,t})\| \|R_{s,t}^y\| d\eta + \int_0^1 \|Dg(y_s + \eta y_{s,t}) - Dg(y_s)\| \|g(y_s)\| \|x_{s,t}\| d\eta \\ &\leq C_g \|R_{s,t}^y\| + \frac{1}{2} C_g^2 \|y_{s,t}\| \|x_{s,t}\|. \end{aligned}$$

Hence, it follows, using p -variation norms and Hölder inequality, that

$$\begin{aligned} \left\| [g(y)]' \right\|_{p\text{-var},[s,t]} &\leq 2C_g^2 \|y\|_{p\text{-var},[s,t]}, \quad \left\| [g(y)]' \right\|_{\infty,[s,t]} \leq C_g^2, \\ \left\| R^{g(y)} \right\|_{q\text{-var},[s,t]} &\leq C_g \|R^y\|_{q\text{-var},[s,t]} + \frac{1}{2}C_g^2 \|x\|_{p\text{-var},[s,t]} \|y\|_{p\text{-var},[s,t]}. \end{aligned} \quad (4.2)$$

As a result, by introducing $\|y, R\|_{p\text{-var},[s,t]} := \|y\|_{p\text{-var},[s,t]} + \|R^y\|_{q\text{-var},[s,t]}$, we obtain

$$\begin{aligned} \|y_{s,t}\| &\leq \left\| \int_s^t g(y_u) dx_u \right\| \\ &\leq \|g(y_s)\| \|x_{s,t}\| + \|Dg(y_s)g(y_s)\| \|\mathbb{X}_{s,t}\| \\ &\quad + C_p \left[\|x\|_{p\text{-var},[s,t]} \left\| R^{g(y)} \right\|_{q\text{-var},[s,t]} + \|\mathbb{X}\|_{q\text{-var},[s,t]} \left\| [g(y)]' \right\|_{p\text{-var},[s,t]} \right] \\ &\leq C_g \|x\|_{p\text{-var},[s,t]} + C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \\ &\quad + C_p \left\{ C_g \|x\|_{p\text{-var},[s,t]} \|R^y\|_{q\text{-var},[s,t]} + \left[\frac{1}{2} \|x\|_{p\text{-var},[s,t]}^2 + 2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \right] C_g^2 \|y\|_{p\text{-var},[s,t]} \right\} \\ &\leq C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} + C_g^2 \|\mathbf{x}\|_{p\text{-var},[s,t]}^2 + 2C_p \left\{ C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \vee C_g^2 \|\mathbf{x}\|_{p\text{-var},[s,t]}^2 \right\} \|y, R\|_{p\text{-var},[s,t]}, \\ &\leq 2 \left\{ C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \vee C_g^2 \|\mathbf{x}\|_{p\text{-var},[s,t]}^2 \right\} \left(1 + C_p \|y, R\|_{p\text{-var},[s,t]} \right), \end{aligned}$$

which yields

$$\|y\|_{p\text{-var},[s,t]} \leq 2 \left\{ C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \vee C_g^2 \|\mathbf{x}\|_{p\text{-var},[s,t]}^2 \right\} \left(1 + C_p \|y, R\|_{p\text{-var},[s,t]} \right). \quad (4.3)$$

The same estimate for R^y is actually included in the above estimate, hence

$$\|R^y\|_{q\text{-var},[s,t]} \leq 2 \left\{ C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \vee C_g^2 \|\mathbf{x}\|_{p\text{-var},[s,t]}^2 \right\} \left(1 + C_p \|y, R\|_{p\text{-var},[s,t]} \right). \quad (4.4)$$

Combining (4.3) and (4.4) gives

$$\|y, R\|_{p\text{-var},[s,t]} \leq 4 \left\{ C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \vee C_g^2 \|\mathbf{x}\|_{p\text{-var},[s,t]}^2 \right\} \left(1 + C_p \|y, R\|_{p\text{-var},[s,t]} \right). \quad (4.5)$$

It implies from (4.5) that $\|y, R\|_{p\text{-var},[s,t]} \leq \frac{1}{C_p}$ whenever $\left\{ C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \vee C_g^2 \|\mathbf{x}\|_{p\text{-var},[s,t]}^2 \right\} \leq \frac{1}{8C_p} < 1$, which yields

$$\|y, R\|_{p\text{-var},[s,t]} \leq \frac{1}{C_p} \quad \text{whenever} \quad \|\mathbf{x}\|_{p\text{-var},[s,t]} \leq \frac{1}{8C_p C_g}.$$

By constructing the sequence of greedy times $\{\tau_k(\frac{1}{8C_p C_g})\}_{k \in \mathbb{N}}$ on interval $[a, b]$, it follows from induction that

$$\begin{aligned} \|y_{\tau_{k+1}}\| &\leq \|y\|_{\infty, [\tau_k, \tau_{k+1}]} \leq \|y\|_{p\text{-var}, [\tau_k, \tau_{k+1}]} \leq \|y_{\tau_k}\| + \|y, R\|_{p\text{-var}, [\tau_k, \tau_{k+1}]} \leq \|y_{\tau_k}\| + \frac{1}{C_p} \\ &\leq \dots \leq \|y_a\| + (k+1) \frac{1}{C_p}, \quad \forall k = 0, \dots, N_{[a,b]}(\mathbf{x}) - 1. \end{aligned}$$

That means

$$\|y\|_{\infty, [a,b]} \leq \|y_a\| + \frac{1}{C_p} N_{[a,b]}(\mathbf{x}) \quad (4.6)$$

On the other hand, it then follows from inequality of p -variation seminorm in [8] that

$$\|y, R\|_{p\text{-var}, [a,b]} \leq N_{[a,b]}^{\frac{p-1}{p}}(\mathbf{x}) \sum_{k=0}^{N_{[a,b]}(\mathbf{x})-1} \|y, R\|_{q\text{-var}, [\tau_k, \tau_{k+1}]} \leq \frac{1}{C_p} N_{[a,b]}^{\frac{2p-1}{p}}(\mathbf{x}). \quad (4.7)$$

Step 2: Next, using (4.6) and (4.7) for any two solutions $y_t(\mathbf{x}, y_a)$ and $\bar{y}_t(\mathbf{x}, \bar{y}_a)$ within the bounded range $\frac{1}{C_p} N_{[a,b]}^{\frac{2p-1}{p}}(\mathbf{x})$, let us consider their difference $z_t = \bar{y}_t - y_t$, which satisfies the integral rough equation

$$z_t = z_a + \int_a^t [g(\bar{y}_s) - g(y_s)] dx_s.$$

As a result, $y'_s = g(y_s)$, $\bar{y}'_s = g(\bar{y}_s)$ and

$$\begin{aligned} & g(\bar{y}_t) - g(y_t) - g(\bar{y}_s) + g(y_s) \\ &= \int_0^1 \left[Dg(\bar{y}_s + \eta \bar{y}_{s,t}) \bar{y}_{s,t} - Dg(y_s + \eta y_{s,t}) y_{s,t} \right] d\eta \\ &= \left[Dg(\bar{y}_s) g(\bar{y}_s) - Dg(y_s) g(y_s) \right] \otimes x_{s,t} + \int_0^1 \left[Dg(\bar{y}_s + \eta \bar{y}_{s,t}) R_{s,t}^{\bar{y}} - Dg(y_s + \eta y_{s,t}) R_{s,t}^y \right] d\eta \\ &\quad + \int_0^1 \left\{ \left[Dg(\bar{y}_s + \eta \bar{y}_{s,t}) - Dg(\bar{y}_s) \right] g(\bar{y}_s) - \left[Dg(y_s + \eta y_{s,t}) - Dg(y_s) \right] g(y_s) \right\} \otimes x_{s,t} d\eta \\ &= \left[Dg(\bar{y}_s) g(\bar{y}_s) - Dg(y_s) g(y_s) \right] \otimes x_{s,t} \\ &\quad + \int_0^1 \left\{ Dg(\bar{y}_s + \eta \bar{y}_{s,t}) \left(R_{s,t}^{\bar{y}} - R_{s,t}^y \right) + \left[Dg(\bar{y}_s + \eta \bar{y}_{s,t}) - Dg(y_s + \eta y_{s,t}) \right] R_{s,t}^y \right\} d\eta \\ &\quad + \int_0^1 \left[Dg(\bar{y}_s + \eta \bar{y}_{s,t}) - Dg(\bar{y}_s) \right] \left[g(\bar{y}_s) - g(y_s) \right] \otimes x_{s,t} d\eta \\ &\quad + \int_0^1 \left[Dg(\bar{y}_s + \eta \bar{y}_{s,t}) - Dg(\bar{y}_s) - Dg(y_s + \eta y_{s,t}) + Dg(y_s) \right] g(y_s) \otimes x_{s,t} d\eta \\ &= \left[Dg(\bar{y}_s) g(\bar{y}_s) - Dg(y_s) g(y_s) \right] \otimes x_{s,t} \\ &\quad + \int_0^1 \left\{ Dg(\bar{y}_s + \eta \bar{y}_{s,t}) R_{s,t}^{\bar{y}-y} + \left[Dg(\bar{y}_s + \eta \bar{y}_{s,t}) - Dg(y_s + \eta y_{s,t}) \right] R_{s,t}^y \right\} d\eta \\ &\quad + \int_0^1 \left[Dg(\bar{y}_s + \eta \bar{y}_{s,t}) - Dg(\bar{y}_s) \right] \left[g(\bar{y}_s) - g(y_s) \right] \otimes x_{s,t} d\eta \\ &\quad + \left(\int_0^1 \int_0^1 D^2 g(\bar{y}_s + \mu \eta \bar{y}_{s,t}) \eta (\bar{y}_{s,t} - y_{s,t}) d\mu d\eta \right) g(y_s) \otimes x_{s,t} \\ &\quad + \left(\int_0^1 \int_0^1 \left[D^2 g(\bar{y}_s + \mu \eta \bar{y}_{s,t}) - D^2 g(y_s + \mu \eta y_{s,t}) \right] d\mu d\eta \right) g(y_s) \otimes x_{s,t}. \end{aligned}$$

This proves $[g(\bar{y}) - g(y)]'_s = Dg(\bar{y}_s) g(\bar{y}_s) - Dg(y_s) g(y_s) = Q(\bar{y}_s) - Q(y_s)$ with $\|Q(\bar{y}_s) - Q(y_s)\| \leq 2C_g^2 \|z_s\|$ and

$$\begin{aligned} \|Q(\bar{y}) - Q(y)\|_{p\text{-var},[s,t]} &\leq C_Q \left(\|z\|_{p\text{-var},[s,t]} + \|z\|_{\infty,[s,t]} \|y\|_{p\text{-var},[s,t]} \right) \\ &\leq 2C_g^2 \left(\|z\|_{p\text{-var},[s,t]} + \|z\|_{\infty,[s,t]} \|y\|_{p\text{-var},[s,t]} \right); \end{aligned}$$

and moreover

$$\begin{aligned} \left\| R^{g(\bar{y})-g(y)} \right\|_{q\text{-var},[s,t]} &\leq C_g \|R^z\|_{q\text{-var},[s,t]} + C_g \|z\|_{\infty,[s,t]} \|R^y\|_{q\text{-var},[s,t]} \\ &\quad + \frac{1}{2} C_g^2 \|x\|_{p\text{-var},[s,t]} \left[\|z\|_{p\text{-var},[s,t]} + \|z\|_{\infty} \left(\|\bar{y}\|_{p\text{-var},[s,t]} + \|y\|_{p\text{-var},[s,t]} \right) \right]. \end{aligned}$$

Using the fact that

$$\|z_{s,t}\| \leq \left\| \int_s^t [g(\bar{y}_u) - g(y_u)] dx_u \right\|$$

$$\begin{aligned} &\leq C_g \|z_s\| \|x\|_{p\text{-var},[s,t]} + 2C_g^2 \|z_s\| \|\mathbb{X}\|_{q\text{-var},[s,t]} \\ &\quad + C_p \left\{ \|x\|_{p\text{-var},[s,t]} \left\| R^{g(\bar{y})-g(y)} \right\|_{q\text{-var},[s,t]} + \|\mathbb{X}\|_{q\text{-var},[s,t]} \left\| [g(\bar{y}) - g(y)]' \right\|_{p\text{-var},[s,t]} \right\}, \end{aligned}$$

we can now estimate

$$\begin{aligned} \|z\|_{p\text{-var},[s,t]} &\leq 2 \left\{ C_g \|x\|_{p\text{-var},[s,t]} \vee C_g^2 \|x\|_{p\text{-var},[s,t]}^2 \right\} \times \\ &\quad \times \left\{ \|z\|_{\infty,[s,t]} \left[1 + C_p (\|\bar{y}, R\|_{p\text{-var},[s,t]} + \|y, R\|_{p\text{-var},[s,t]}) \right] + C_p \|z, R\|_{p\text{-var},[s,t]} \right\} \\ &\leq 2C_p \left\{ C_g \|x\|_{p\text{-var},[s,t]} \vee C_g^2 \|x\|_{p\text{-var},[s,t]}^2 \right\} \times \\ &\quad \times \left(1 + \|\bar{y}, R\|_{p\text{-var},[s,t]} + \|y, R\|_{p\text{-var},[s,t]} \right) \left(\|z_s\| + \|z, R\|_{p\text{-var},[s,t]} \right). \end{aligned}$$

The similar estimate for $\|R^z\|_{q\text{-var},[s,t]}$ is already included in the above computation. Therefore by combining with (4.7), we obtain

$$\begin{aligned} &\|z, R\|_{p\text{-var},[s,t]} \\ &\leq 4C_p \left\{ C_g \|x\|_{p\text{-var},[s,t]} \vee C_g^2 \|x\|_{p\text{-var},[s,t]}^2 \right\} \times \\ &\quad \times \left(1 + \|\bar{y}, R\|_{p\text{-var},[s,t]} + \|y, R\|_{p\text{-var},[s,t]} \right) \left(\|z_s\| + \|z, R\|_{p\text{-var},[s,t]} \right) \\ &\leq 4C_p \left\{ C_g \|x\|_{p\text{-var},[s,t]} \vee C_g^2 \|x\|_{p\text{-var},[s,t]}^2 \right\} \left(1 + \frac{2}{C_p} N^{\frac{2p-1}{p}} \frac{1}{8C_p C_g}, [a,b]}(\mathbf{x}) \right) \left(\|z_s\| + \|z, R\|_{p\text{-var},[a,b]} \right), \end{aligned}$$

which follows that

$$\|z, R\|_{p\text{-var},[s,t]} \leq \|z_s\| \quad \text{whenever} \quad 8C_p C_g \left(1 + \frac{2}{C_p} N^{\frac{2p-1}{p}} \frac{1}{8C_p C_g}, [a,b]}(\mathbf{x}) \right) \|x\|_{p\text{-var},[s,t]} \leq 1.$$

Therefore, (2.12) is followed directly from the usage of greedy times (2.13), which is similar to (4.6) and (4.7). \square

Proof: [**Proposition 2.2**] The proof is divided into several steps.

Step 1: First, for a fixed solution $y_t(\mathbf{x}, y_a)$ on a given interval $[a, b]$, we need to prove the existence and uniqueness of the solution of the linearized rough differential equation (2.14), which has the time dependent coefficient $\Sigma_t := Dg(y_t)$. To do that, we simply follow Gubinelli's method by considering the Ito-Lyons map $H_t = \xi_a + \int_a^t \Sigma_s \xi_s dx_s$ on the set $\mathcal{D}_x^{2\beta}([a, b], \xi_a, \Sigma_a \xi_a)$ of controlled paths ξ_t such that ξ_a is fixed, $\xi'_a = \Sigma_a \xi_a$. Note that $\Sigma_t = Dg(y_t)$ is also controlled by x with

$$\Sigma_{s,t} = \int_0^1 D^2 g(y_s + \eta y_{s,t})(g(y_s) \otimes x_{s,t} + R_{s,t}^y) d\eta,$$

thus

$$\Sigma'_s = D_g^2(y_s) g(y_s), \quad \|R_{s,t}^\Sigma\| \leq C_g \mathbb{R}_{s,t}^y + \frac{1}{2} C_g^2 \|y_{s,t}\| \|x_{s,t}\|.$$

As a result $\Sigma_t \xi_t$ is also controlled by x with $[\Sigma \cdot \xi]'_s = \Sigma'_s \xi_s + \Sigma_s \xi'_s$ and

$$\|R_{s,t}^{\Sigma \cdot \xi}\| \leq \|\Sigma_{s,t}\| \|\xi_{s,t}\| + \|\xi_s\| \|R_{s,t}^\Sigma\| + \|\Sigma_s\| \|R_{s,t}^\xi\|.$$

It then enable to estimate

$$\begin{aligned} &\|H_{s,t} - \Sigma_s \xi_s \otimes x_{s,t} + [\Sigma'_s \xi_s + \Sigma_s \xi'_s] \otimes \mathbb{X}_{s,t}\| \\ &= \left\| \int_s^t \Sigma_u \xi_u dx_u - \Sigma_s \xi_s \otimes x_{s,t} + [\Sigma'_s \xi_s + \Sigma_s \xi'_s] \otimes \mathbb{X}_{s,t} \right\| \\ &\quad + C_\beta (t-s)^{3\beta} \left(\|x\|_{\beta,[s,t]} \left\| R^{\Sigma \cdot \xi} \right\|_{2\beta,[s,t]} + \|\mathbb{X}\|_{2\beta,[s,t]} \left\| [\Sigma \cdot \xi]' \right\|_{\beta,[s,t]} \right) \end{aligned}$$

where

$$\begin{aligned}
\|[\Sigma \cdot \xi]'\|_\beta &\leq \|\Sigma'\|_\infty \|\xi\|_\beta + \|\Sigma'\|_\beta \|\xi\|_\infty + \|\Sigma\|_\infty \|\xi'\|_\beta + \|\Sigma\|_\beta \|\xi'\|_\infty, \\
\|R^{\Sigma \cdot \xi}\|_{2\beta} &\leq \|\Sigma\|_\beta \|\xi\|_\beta + \|R^\Sigma\|_{2\beta} \|\xi\|_\infty + \|\Sigma\|_\infty \|R^\xi\|_{2\beta} \quad \text{with} \\
\|\xi'\| &\leq \|\xi'_a\| + (t-a)^\beta \|\xi'\|_\beta \leq \|\Sigma\|_\infty \|\xi_a\| + (t-a)^\beta \|\xi, \xi'\|_{2\beta}, \\
\|\xi\|_\beta &\leq \|\xi'\|_\infty \|x\|_\beta + (t-a)^\beta \|R^\xi\|_{2\beta} \leq \|\Sigma\|_\infty \|x\|_\beta \|\xi_a\| + (\|x\|_\beta \vee 1) (t-a)^\beta \|\xi, \xi'\|_{2\beta}, \\
\|\xi\|_\infty &\leq \|\xi_a\| + (t-a)^\beta \|\xi\|_\beta \leq \|\xi_a\| \left(1 + \|\Sigma\|_\infty (t-a)^\beta \|x\|_\beta\right) + (t-a)^{2\beta} (\|x\|_\beta \vee 1) \|\xi, \xi'\|_{2\beta}.
\end{aligned}$$

A direct computation then shows that

$$\begin{aligned}
\|[\Sigma \cdot \xi]'\|_\beta &\leq \left\{ \|\Sigma\|_\infty + (t-a)^\beta \|\Sigma\|_\beta + \|\Sigma'\|_\infty (t-a)^\beta (\|x\|_\beta \vee 1) + \|\Sigma'\|_\beta (t-a)^{2\beta} (\|x\|_\beta \vee 1) \right\} \|\xi, \xi'\|_{2\beta} \\
&\quad + \left\{ \|\Sigma\|_\beta \|\Sigma\|_\infty + \|\Sigma'\|_\infty \|\Sigma\|_\infty \|x\|_\beta + \|\Sigma'\|_\beta \left(1 + \|\Sigma\|_\infty (t-a)^\beta \|x\|_\beta\right) \right\} \|\xi_a\| \\
\|R^{\Sigma \cdot \xi}\|_{2\beta} &\leq \left\{ \|\Sigma\|_\infty + \|\Sigma\|_\beta (t-a)^\beta (\|x\|_\beta \vee 1) + \|R^\Sigma\|_{2\beta} (t-a)^{2\beta} (\|x\|_\beta \vee 1) \right\} \|\xi, \xi'\|_{2\beta} \\
&\quad + \left\{ \|\Sigma\|_\beta \|\Sigma\|_\infty \|x\|_\beta + \|R^\Sigma\|_{2\beta} \left(1 + \|\Sigma\|_\infty (t-a)^\beta \|x\|_\beta\right) \right\} \|\xi_a\|.
\end{aligned}$$

Hence

$$\begin{aligned}
\|R^H\|_{2\beta} &\leq \left(\|\Sigma'\|_\infty \|\xi\|_\infty + \|\Sigma\|_\infty \|\xi'\|_\infty \right) \|\mathbb{X}\|_{2\beta} + C_\beta (t-a)^\beta \left(\|x\|_\beta \|R^{\Sigma \cdot \xi}\|_{2\beta} + \|\mathbb{X}\|_{2\beta} \|[\Sigma \cdot \xi]'\|_\beta \right) \\
\|H'\|_\beta &\leq \|\Sigma\|_\infty \|\xi\|_\beta + \|\Sigma\|_\beta \|\xi\|_\infty \\
&\leq \left\{ \|\Sigma\|_\infty (t-a)^\beta (\|x\|_\beta \vee 1) + \|\Sigma\|_\beta (t-a)^{2\beta} (\|x\|_\beta \vee 1) \right\} \|\xi, \xi'\|_{2\beta} \\
&\quad + \left\{ \|\Sigma\|_\infty^2 \|x\|_\beta + \|\Sigma\|_\beta \left[1 + \|\Sigma\|_\infty (t-a)^\beta \|x\|_\beta\right] \right\} \|\xi_a\|.
\end{aligned}$$

Combining everything together, we have just showed that there exists constants

$$M_1 = M_1(\Sigma, [a, b], x, \mathbb{X}), \quad M_2 = M_2(\Sigma, [a, b], x, \mathbb{X})$$

such that

$$\|H'\|_\beta + \|R^H\|_{2\beta} \leq M_2 \|\xi_a\| + M_1 \left[(t-a)^\beta + \|x\|_\beta + \|\mathbb{X}\|_{2\beta} \right] \|\xi, \xi'\|_{2\beta}.$$

This implies that on every interval $[t_k, t_{k+1}]$ of the greedy times

$$t_0 = a, \quad t_{k+1} = \inf\{t > t_k : M_1 \left[(t-t_k)^\beta + \|x\|_{\beta, [t_k, t]} + \|\mathbb{X}\|_{2\beta, [t_k, t]} = \frac{1}{2} \right] \wedge b,$$

the Ito-Lyons map is a contraction from the set $\left\{ \mathcal{D}_x^{2\beta}([a, b], \xi_a, \Sigma_a \xi_a) : \|\xi, \xi'\|_{2\beta, [t_k, t_{k+1}]} \leq 2M_2 \|\xi_{t_k}\| \right\}$ into itself, hence there exists a unique solution of (2.14) on every interval $[t_k, t_{k+1}]$. The concatenation of solutions on intervals $[t_k, t_{k+1}]$ then proves the existence and uniqueness of the solution of (2.14) on $[a, b]$.

Step 2: Denote by $\Phi(t, x, z_a)$ the solution matrix of the linearized system (2.14), then $\xi = \Phi(t, x, z_a)(\bar{z}_a - z_a)$ is the solution of (2.14) given initial point $\xi_a = \bar{z}_a - z_a$. Assign $r_t := \bar{z}_t - z_t - \xi_t$, then $r_a = 0$ and

$$\begin{aligned}
r_t &= \int_a^t \left[\int_0^1 D_z g(z_s + \eta(\bar{z}_s - z_s)) - D_z g(z_s) \right] (\bar{z}_s - z_s) d\eta dx_s + \int_a^t D_z g(z_s) r_s dx_s, \\
&= e_{a,t} + \int_a^t D_z g(z_s) r_s dx_s, \quad \forall t \in [a, b],
\end{aligned} \tag{4.8}$$

where

$$e_{a,t} = \int_a^t \int_0^1 \left[D_z g(z_s + \eta(\bar{z}_s - z_s)) - D_z g(z_s) \right] (\bar{z}_s - z_s) d\eta dx_s$$

and e is also controlled by x with $e_a = 0$. We are going to estimate $\|r\|_{\infty,[a,b]}$ and $\|r, R\|_{p\text{-var},[a,b]}$ through $\|e'\|_{\infty,[a,b]}$, $\|e'\|_{p\text{-var},[a,b]}$, $\|R^e\|_{q\text{-var},[a,b]}$. First observe that

$$r_{s,t} = e_{s,t} + \int_s^t Dg(y_u) r_u dx_u = e' \otimes x_{s,t} + R_{s,t}^e + \int_s^t Dg(y_u) r_u dx_u,$$

which yields $r'_s = e'_s + Dg(y_s)r_s$ and

$$\|R_{s,t}^r\| \leq \|R_{s,t}^e\| + \|[Dg(y)r]'\|_s \|\mathbb{X}_{s,t}\| + C_p \left(\|x\|_p \left\| R^{Dg(y)r} \right\|_q + \|\mathbb{X}\|_q \|[Dg(y)r]'\|_p \right). \quad (4.9)$$

A direct computation shows that

$$\begin{aligned} & \left\| Dg(y_t)r_t - Dg(y_s)r_s - \left[D_g^2(y_s)g(y_s)r_s + Dg(y_s)r'_s \right] \otimes x_{s,t} \right\| \\ & \leq \left\| \int_0^1 D^2g(y_s + \eta y_{s,t}) R_{s,t}^y r_s d\eta \right\| + \frac{1}{2} C_g^2 \|y_{s,t}\| \|r_s\| \|x_{s,t}\| + \|Dg(y_s)R_{s,t}^y\| + C_g \|y_{s,t}\| \|r_{s,t}\|, \end{aligned}$$

which yields $[Dg(y)r]'_s = D^2g(y_s)[g(y_s), r_s] + Dg(y_s)r'_s$ and

$$\begin{aligned} \|R_{s,t}^{D(y)r}\| & \leq C_g \|R_{s,t}^y\| \|r\|_{\infty} + \frac{1}{2} C_g^2 \|r\|_{\infty} \|y_{s,t}\| \|x_{s,t}\| + C_g \|R_{s,t}^r\| + C_g \|y_{s,t}\| \|r_{s,t}\| \\ \Rightarrow \left\| R^{Dg(y)r} \right\|_q & \leq C_g \|r\|_{\infty} \|R^y\|_q + \frac{1}{2} C_g^2 \|r\|_{\infty} \|y\|_p \|x\|_p + C_g \|R^r\|_q + C_g \|y\|_p \|r\|_p. \end{aligned}$$

Similarly, we can show that

$$\begin{aligned} \|[Dg(y)r]'\|_{\infty} & \leq C_g^2 \|r\|_{\infty} + C_g \|r'\|_{\infty} \\ \|[Dg(y)r]'\|_p & \leq 2C_g^2 \|r\|_{\infty} \|y\|_p + C_g^2 \|r\|_p + C_g \|r'\|_p + C_g \|r'\|_{\infty} \|y\|_p. \end{aligned}$$

Combining all the above estimates into (4.9), we obtain

$$\begin{aligned} \|R^r\|_q & \leq C_p C_g \|x\|_p \|R^r\|_q \\ & + \|r\|_p \left\{ 2C_g^2 \|\mathbb{X}\|_q + C_p \|x\|_p \left[C_g \|R^y\|_q + \frac{1}{2} C_g^2 \|x\|_p \|y\|_p + C_g \|y\|_p \right] + C_g \|\mathbb{X}\|_q (4C_g^2 \|y\|_p + 2C_g^2) \right\} \\ & + \|r_s\| \left\{ 2C_g^2 \|\mathbb{X}\|_q + C_p \|x\|_p \left[C_g \|R^y\|_q + \frac{1}{2} C_g^2 \|x\|_p \|y\|_p \right] + 4C_p C_g^2 \|\mathbb{X}\|_q \right\} \\ & + \|R^e\|_q + C_g \|\mathbb{X}\|_q \|e'\|_{\infty} + C_p \|\mathbb{X}\|_q (C_g \|e'\|_p + C_g \|y\|_p \|e'\|_{\infty}) =: \bar{R}, \end{aligned}$$

and similarly

$$\|r\|_p \leq \|e'\|_{\infty} \|x\|_p + C_g \|r\|_{\infty} \|x\|_p + \bar{R} \leq \|e'\|_{\infty} \|x\|_p + C_g \|x\|_p (\|r_s\| + \|r\|_p) + \bar{R}.$$

Therefore, taking into account (4.7) we have just proved that there exists a constant

$$M = M(p, [a, b], \|\mathbf{x}\|_{p\text{-var},[a,b]}) > 1$$

such that

$$\begin{aligned} \|r, R\|_{p\text{-var},[s,t]} & \leq M \left(C_g \|x\|_{p\text{-var},[s,t]} + C_g^2 \|\mathbb{X}\|_{q\text{-var},[s,t]} \right) \left(\|r_s\| + \|r, R\|_{p\text{-var},[s,t]} \right) \\ & + M \left(\|e'\|_{\infty,[a,b]} + \|e'\|_{p\text{-var},[a,b]} + \|R^e\|_{q\text{-var},[a,b]} \right) \\ & \leq 2M \left(C_g \|\mathbf{x}\|_{p\text{-var},[s,t]} \vee C_g^2 \|\mathbf{x}\|_{q\text{-var},[s,t]}^2 \right) \left(\|r_s\| + \|r, R\|_{p\text{-var},[s,t]} \right) \\ & + M \left(\|e'\|_{\infty,[a,b]} + \|e'\|_{p\text{-var},[a,b]} + \|R^e\|_{q\text{-var},[a,b]} \right), \end{aligned}$$

which implies

$$\|r, R\|_{p\text{-var},[s,t]} \leq \|r_s\| + 2M \left(\|e'\|_{\infty,[a,b]} + \|e'\|_{p\text{-var},[a,b]} + \|R^e\|_{q\text{-var},[a,b]} \right)$$

whenever $2MC_g \|x\|_{p\text{-var},[s,t]} \leq \frac{1}{2}$. Similar estimates to (4.6) and (4.7), using the sequence of greedy times $\{\tau_k(\frac{1}{4MC_g})\}_{k \in \mathbb{N}}$, lead to

$$\begin{aligned} & \|r\|_{\infty,[a,b]} \vee \|r, R\|_{p\text{-var},[s,t]} \\ & \leq N_{\frac{1}{4MC_g},[a,b]}(\mathbf{x})^{\frac{p-1}{p}} \left\{ \|r_a\| + 2M \left(\|e'\|_{\infty,[a,b]} + \|e'\|_{p\text{-var},[a,b]} + \|R^e\|_{q\text{-var},[a,b]} \right) \right\} e^{(\log 2)N_{\frac{1}{4MC_g},[a,b]}(\mathbf{x})} \\ & \leq 2M \left(\|e'\|_{\infty,[a,b]} + \|e'\|_{p\text{-var},[a,b]} + \|R^e\|_{q\text{-var},[a,b]} \right) e^{(1+\log 2)N_{\frac{1}{4MC_g},[a,b]}(\mathbf{x})}, \end{aligned} \quad (4.10)$$

where we use the fact that $r_a = 0$.

Step 3: From (4.8), it follows that

$$e'_s = \int_0^1 [Dg(y_s + \eta z_s) - Dg(y_s)] z_s d\eta = \int_0^1 \int_0^1 D^2g\left((1 - \mu\eta)y_s + \mu\eta\bar{y}_s\right) [z_s, z_s] \eta d\mu d\eta = \bar{e}_s,$$

is controlled by x . As a result $\|e'_s\| \leq \frac{1}{2}C_g \|z_s\|^2$ thus

$$\|e'\|_{\infty,[a,b]} \leq \frac{1}{2}C_g \|z\|_{\infty,[a,b]}^2, \quad (4.11)$$

$$\|e'\|_{p\text{-var},[a,b]} \leq C_g \left(\|y\|_{p\text{-var},[a,b]} \vee \|\bar{y}\|_{p\text{-var},[a,b]} \right) \|z\|_{\infty,[a,b]}^2 + 2C_g \|z\|_{\infty,[a,b]} \|z\|_{p\text{-var},[a,b]}.$$

On the other hand,

$$\|R_{s,t}^e\| \leq \|e'\|_{\infty} \|\mathbb{X}_{s,t}\| + C_p \left(\|x\|_{p\text{-var},[a,b]} \|R^{\bar{e}}\|_{q\text{-var},[a,b]} + \|\mathbb{X}\|_{q\text{-var},[a,b]} \|e'\|_{p\text{-var},[a,b]} \right)$$

thus $\|R^e\|_{q\text{-var},[a,b]} \leq \|e'\|_{\infty,[a,b]} \|\mathbb{X}\|_{q\text{-var},[a,b]}$

$$+ C_p \left(\|x\|_{p\text{-var},[a,b]} \|R^{\bar{e}}\|_{q\text{-var},[a,b]} + \|\mathbb{X}\|_{q\text{-var},[a,b]} \|e'\|_{p\text{-var},[a,b]} \right), \quad (4.12)$$

where a direct computation shows that

$$\begin{aligned} \bar{e}'_s &= \int_0^1 \int_0^1 \left\{ D^3g\left((1 - \mu\eta)y_s + \mu\eta\bar{y}_s\right) [(1 - \mu\eta)y'_s + \mu\eta\bar{y}'_s, z_s, z_s] \right. \\ & \quad \left. + 2D^2g\left((1 - \mu\eta)y_s + \mu\eta\bar{y}_s\right) [z'_s, z_s] \right\} \eta d\mu d\eta, \\ R_{s,t}^{\bar{e}} &= \int_0^1 \int_0^1 \left\{ R_{s,t}^{D^2g(\bar{y},y)} [z_t, z_t] + D^2g(\bar{y}, y) \left([z_s, R_{s,t}^z] + [z'_s x_{s,t}, z'_s x_{s,t} + R_{s,t}^z] + [R_{s,t}^z, z_t] \right) \right. \\ & \quad \left. + [D^2g(\bar{y}, y)]'_s x_{s,t} \left([z_s + z_t, z'_s + R_{s,t}^z] \right) \right\} \eta d\mu d\eta. \end{aligned}$$

We therefore can show that there exists a generic constant α such that

$$\begin{aligned} & \|e'\|_{\infty,[a,b]} \vee \|e'\|_{p\text{-var},[a,b]} \vee \|R^{\bar{e}}\|_{q\text{-var},[a,b]} \\ & \leq \alpha \left(\|z\|_{\infty,[a,b]} + \|z, R\|_{p\text{-var},[a,b]} + \|z'\|_{\infty,[a,b]} + \|z'\|_{p\text{-var},[a,b]} \right)^2. \end{aligned} \quad (4.13)$$

By replacing (4.11), (4.12), (4.13) into (4.10), and using (2.12), we derive that there exists a generic constant such that

$$\|\bar{y}(\mathbf{x}, \bar{y}_a) - y(\mathbf{x}, y_a) - \xi(\mathbf{x}, \bar{y}_a - y_a)\|_{\infty,[a,b]} \leq \alpha \|\bar{y}_a - y_a\|^2.$$

This, combined with the linearity of ξ w.r.t. $\bar{y}_a - y_a$, shows the differentiability of $y_t(\mathbf{x}, y_a)$ w.r.t. y_a . □

Acknowledgments

The author would like to thank Vietnam Institute for Advanced Studies in Mathematics (VIASM) for financial support during three month research stay at the institute in 2019.

References

- [1] H. Amann. *Ordinary Differential Equations: An Introduction to Nonlinear Analysis*. Walter de Gruyter, Berlin . New York, 1990.
- [2] L. Arnold. *Random Dynamical Systems*. Springer, Berlin Heidelberg New York, 1998.
- [3] I. Bailleul, S. Riedel, M. Scheutzow. *Random dynamical systems, rough paths and rough flows*. J. Differential Equations, Vol. **262**, (2017), 5792–5823.
- [4] T. Cass, C. Litterer, T. Lyons. *Integrability and tail estimates for Gaussian rough differential equations*. Annals of Probability, Vol. **14**, No. 4, (2013), 3026–3050.
- [5] L.Coutin. *Rough paths via sewing lemma*. ESAIM: Probability and Statistics., **16**, (2012), 479–526.
- [6] H. Crauel, P. Kloeden, *Nonautonomous and random attractors*. Jahresber Dtsch. Math-Ver. **117** (2015), 173–206.
- [7] L. H. Duc. *Stability theory for Gaussian rough differential equations. Part I*. Preprint.
- [8] L. H. Duc. *Stability theory for Gaussian rough differential equations. Part II*. Preprint.
- [9] L. H. Duc, M. J. Garrido-Atienza, A. Neuenkirch, B. Schmalfuß. *Exponential stability of stochastic evolution equations driven by small fractional Brownian motion with Hurst parameter in $(\frac{1}{2}, 1)$* . Journal of Differential Equations, 264 (2018), 1119-1145.
- [10] L. H. Duc, P. T. Hong. *Asymptotic dynamics for Young differential equations: a unified approach*. Preprint. ArXiv: 1905.04945.
- [11] P. Friz, M. Hairer. *A course on rough path with an introduction to regularity structure*. Universitext, Vol. **XIV**, Springer, Berlin, 2014.
- [12] P. Friz, N. Victoir. *Multidimensional stochastic processes as rough paths: theory and applications*. Cambridge Studies in Advanced Mathematics, 120. Cambridge University Press, Cambridge, 2010.
- [13] M. Garrido-Atienza, B. Maslowski, B. Schmalfuß. *Random attractors for stochastic equations driven by a fractional Brownian motion*. International Journal of Bifurcation and Chaos, Vol. 20, No. 9 (2010) 27612782.
- [14] M. Garrido-Atienza, A. Neuenkirch, B. Schmalfuß. *Asymptotic stability of differential equations driven by Hölder-continuous paths* J. Dyn. Diff. Equat., (2018), in press.
- [15] M. Garrido-Atienza, B. Schmalfuss. *Local Stability of Differential Equations Driven by Hölder-Continuous Paths with Hölder Index in $(\frac{1}{3}, \frac{1}{2})$* . SIAM J. Appl. Dyn. Syst. Vol. **17**, No. **3**, (2018), 2352–2380.
- [16] M. Gubinelli. *Controlling rough paths*. J. Funtional Analysis, **216** (1), (2004), 86–140.
- [17] R. Khasminskii. *Stochastic stability of differential equations*. Springer, Vol. 66, 2011.

- [18] T. Lyons. *Differential equations driven by rough signals*. Rev. Mat. Iberoam., Vol. **14** (2), (1998), 215–310.
- [19] T. Lyons, M. Caruana, Th. Lévy. *Differential equations driven by rough paths*. Lecture Notes in Mathematics, Vol. **1908**, Springer, Berlin 2007.
- [20] B. Mandelbrot, J. van Ness. *Fractional Brownian motion, fractional noises and applications*. SIAM Review, **4**, No. 10, (1968), 422–437.
- [21] X. Mao, *Stochastic differential equations and applications*. Elsevier, 2007.
- [22] I. Nourdin. *Selected aspects of fractional Brownian motion*. Bocconi University Press, Springer, 2012.
- [23] S. Riedel, M. Scheutzow. *Rough differential equations with unbounded drift terms*. J. Differential Equations, Vol. **262**, (2017), 283–312.
- [24] H. J. Sussmann. On the gap between deterministic and stochastic ordinary differential equations. *Ann. Probability*. **6**, No. 1, (1978), 19–41.
- [25] L.C. Young. *An integration of Hölder type, connected with Stieltjes integration*. Acta Math. **67**, (1936), 251–282.